Attorney's Docket No.: 13425-042001 / 00298-US



## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Sven Enerbäck et al.

Art Unit : 1645

Serial No.: 09/9

09/963,285

Examiner:

Filed

September 26, 2001

Title

: PROMOTER SEQUENCES

Commissioner for Patents Washington, D.C. 20231

# TRANSMITTAL OF CERTIFIED PRIORITY DOCUMENT UNDER 35 USC §119

In accordance with the provisions of 35 U.S.C. §119, applicants hereby claim priority of Swedish Patent application No. 0004102-0, filed November 9, 2000. A certified copy of the application is submitted herewith. As the priority application is in the English language, all of the requirements of §119 have been met.

Please apply any charges or credits to Deposit Account No. 06-1050, referencing Attorney Docket No. 13425-042001.

Respectfully submitted,

Date: Date: Ld, dOL

Jack Brennan Reg. No. 47,443

Fish & Richardson P.C. 225 Franklin Street

Boston, Massachusetts 02110-2804

Telephone: (617) 542-5070 Facsimile: (617) 542-8906

20376379.doc

## CERTIFICATE OF MAILING BY FIRST CLASS MAIL

I hereby certify under 37 CFR §1.8(a) that this correspondence is being deposited with the United States Postal Service as first class mail with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

Date of Deposit

Signature

Jarlene J. Morin

Typed or Printed Name of Person Signing Certificate



Intyg C rtificat



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

- (71) Sökande Biovitrum AB, Stockholm SE Applicant (s)
- (21) Patentansökningsnummer 0004102-0 Patent application number
- (86) Ingivningsdatum 2000-11-09 Date of filing

Stockholm, 2001-11-16

För Patent- och registreringsverket For the Patent- and Registration Office

Christina Vängborg

Avgift Fee

170:-

15

20

25

30

ĭ

## PROMOTER SEQUENCES II

### **TECHNICAL FIELD**

The present invention relates an isolated promoter region of the mammalian transcription factor FOXC2. The invention also relates to screening methods for agents modulating the expression of FOXC2 and thereby being potentially useful for the treatment of medical conditions related to obesity. The invention further relates to a previously unknown variant of the human FOXC2 gene, derived via the use of an alternative promoter, which produces an additional exon that generates a distinct open reading frame via splicing. The alternative gene encodes a variant of the FOXC2 transcription factor, which is lacking a part of the DNA-binding domain and consequently has a potential regulatory function.

#### **BACKGROUND ART**

More than half of the men and women in the United States, 30 years of age and older, are now considered overweight, and nearly one-quarter are clinically obese. This high prevalence has led to increases in the medical conditions that often accompany obesity, especially non-insulin dependent diabetes mellitus (NIDDM), hypertension, cardiovascular disorders, and certain cancers. Obesity results from a chronic imbalance between energy intake (feeding) and energy expenditure. To better understand the mechanisms that lead to obesity and to develop strategies in certain patient populations to control obesity, there is a need to develop a better underlying knowledge of the molecular events that regulate the differentiation of preadipocytes and stem cells to adipocytes, the major component of adipose tissue.

The helix-loop-helix (HLH) family of transcriptional regulatory proteins are key players in a wide array of developmental processes (for a review, see Massari & Murre (2000) Mol. Cell. Biol. 20: 429-440). Over 240 HLH proteins have been identified to date in organisms ranging from the yeast *Saccharomyces cerevisiae* to humans. Studies in *Xenopus laevis*, *Drosophila melanogaster*, and mice have convincingly demonstrated that HLH proteins are intimately involved in developmental events such as cellular differentiation, lineage

Ĩ

commitment, and sex determination. In multicellular organisms, HLH factors are required for a multitude of important developmental processes, including neurogenesis, myogenesis, hematopoiesis, and pancreatic development.

The winged helix / forkhead class of transcription factors is characterized by a 100-amino acid, monomeric DNA-binding domain. X-ray crystallography of the forkhead domain from HNF-3γ has revealed a three-dimensional structure, the "winged helix", in which two loops (wings) are connected on the C-terminal side of the helix-loop-helix (for reviews, see Brennan, R.G. (1993) Cell 74: 773-776; and Lai, E. et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90: 10421-10423).

The isolation of the mouse mesenchyme forkhead-1 (MFH-1) and the corresponding human (*FKHL14*) chromosomal genes is disclosed by Miura, N. et al. (1993) FEBS letters 326: 171-176; and (1997) Genomics 41: 489-492. The nucleotide sequences of the mouse MFH-1 gene and the human *FKHL14* gene have been deposited with the EMBL/GenBank Data Libraries under accession Nos. Y08222 (SEQ ID NO: 5) and Y08223 (SEQ ID NO: 8), respectively. A corresponding gene has been identified in *Gallus gallus* (GenBank accession numbers U37273 and U95823).

The International Patent Application WO 98/54216 discloses a gene encoding a Forkhead-Related Activator (FREAC)-11 (also known as S12), which is identical with the polypeptide encoded by the human FKHL14 gene disclosed by Miura, supra. This transcription factor is expressed in adipose tissue and involved in lipid metabolism and adipocyte differentiation (cf. Swedish patent application No. 0000531-4, filed February 18, 2000).

The nomenclature for the winged helix / forkhead transcription factors has been standardized and Fox (Forkhead Box) has been adopted as the unified symbol (Kaestner et al. (2000) Genes & Development 14: 142-146; see also http://www.biology.pomona.edu/fox). It has been agreed that the genes previously designated MFH-1 and FKHL14 (as well as FREAC-11 and S12) should be designated FOXC2.

20

25

30

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the general structure of the human FOXC2 gene.

Figure 2 illustrates the results from phylogenetic footprinting experiments. Shown is the fraction conserved (1.0 = 100%) between mouse FoxC2 and human FOXC2 sequences in the alignment generated with Clustal. Solid (bold) line indicates the fraction of the human sequence which is identical to the mouse within a 200 bp "window" over the human sequence in the alignment. The weak (dotted) line is set to -0.05 when the sliding window contains human exon sequence and to -0.1 when the window is entirely composed of exon sequence. Regions containing local maxima or exceeding a conservation fraction of 0.7 are likely to be functional and are classified as "predicted regulatory regions".

Figure 3 illustrates the predicted "enhancer" region in the human *FOXC2* gene. Underlined sequences indicate likely transcription factor binding sites. Boxed sequence indicates exon sequence.

Splice = sequence predicted as splice site in the alternatively spliced gene;

E-box-like = sequence resembling the "E-box" motif CANNTG known as a target for DNA binding proteins containing a helix-loop-helix domain (often associated with the activation of cell-type specific gene transcription during tissue differentiation; see Massari & Murre (2000) Mol. Cell. Biol. 20: 429-440)

Forkhead-like = sequence resembling binding site for the winged helix / forkhead class of transcription factors;

Ets-like = sequence resembling consensus binding site for ETS-domain transcription factor family (see Sharrocks et al. (1997) Int. J. Biochem. Cell Biol. 29, 1371-1387).

Figure 4 illustrates the predicted "promoter" region in the human *FOXC2* gene. Underlined sequence indicates exon sequences. Boxed sequences indicate conserved block (potential transcription factor binding sites).

10

25

30

1

#### **DESCRIPTION OF THE INVENTION**

According to the present invention, the partially known sequence (SEQ ID NO: 8) of human FOXC2 gene has been extended. In the previously unknown region of the gene, differentially conserved regions, consistent with regulatory function, have been identified. Further, an alternative transcript has been identified, which includes the use of at least two exons. The putative regulatory enhancer is immediately adjacent to the newly discovered alternative exon, suggesting that it may play a role in the alternative selection of transcript classes.

Modulation of the *FOXC2* regulation is expected to have therapeutic value in type II diabetes; obesity, hypercholesterolemia, and other cardiovascular diseases or dyslipidemias.

- 15 Consequently, in a first aspect this invention provides a human FOXC2 promoter region comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 1250 to 2235, such as positions 1250 to 1749 or positions 1692 to 1703, in SEQ ID NO: 1, or a fragment thereof exhibiting *FOXC2* promoter activity;
- 20 (b) the complementary strand of (a); and
  - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).

Another aspect of the invention is a recombinant construct comprising the human FOXC2 promoter region as defined above. In the said recombinant construct, the human FOXC2 promoter region can be operably linked to a gene encoding a detectable product, such as the human FOXC2 gene, or a reporter gene. The term "operably linked" as used herein means functionally fusing a promoter with a structural gene in the proper frame to express the structural gene under control of the promoter. As used herein, the term "reporter gene" means a gene encoding a gene product that can be identified using simple, inexpensive methods or reagents and that can be operably linked to the human FOXC2 promoter region or an active fragment thereof. Reporter genes such as, for example, a luciferase,  $\beta$ -galactosidase, alkaline phosphatase, or green fluorescent protein reporter gene, can be used

G.

to determine transcriptional activity in screening assays according to the invention (see, for example, Goeddel (ed.), Methods Enzymol., Vol. 185, San Diego: Academic Press, Inc. (1990); see also Sambrook, *supra*).

The invention also provides a vector comprising the recombinant construct as defined above, as well as a host cell stably transformed with such a vector, or generally with the recombinant construct according to the invention. The term "vector" refers to any carrier of exogenous DNA that is useful for transferring the DNA to a host cell for replication and/or appropriate expression of the exogenous DNA by the host cell.

10

15

20

25

30

In another aspect, the invention provides a method for identification of an agent regulating FOXC2 promoter activity, said method comprising the steps: (i) contacting a candidate agent with a human FOXC2 promoter region as defined above; and (ii) determining whether said candidate agent modulates expression of the FOXC2 gene, such modulation being indicative for an agent capable of regulating FOXC2 promoter activity. As used herein, the term "agent" means a biological or chemical compound such as a simple or complex organic molecule, a peptide, a protein or an oligonucleotide.

A transfection assay can be a particularly useful screening assay for identifying an effective agent modulating and/or regulating FOXC2 promoter activity. In a transfection assay, a nucleic acid containing a gene, e.g. a reporter gene, operably linked to a human FOXC2 promoter or an active fragment thereof, is transfected into the desired cell type. A test level of reporter gene expression is assayed in the presence of a candidate agent and compared to a control level of expression. An effective agent is identified as an agent that results in a test level of expression that is different than a control level of reporter gene expression, which is the level of expression determined in the absence of the agent.

Methods for transfecting cells and a variety of convenient reporter genes are well known in the art (see, for example, Goeddel (ed.), Methods Enzymol., Vol. 185, San Diego:

Academic Press, Inc. (1990); see also Sambrook, supra). Consequently, the said method could e.g. comprising assaying reporter gene expression in a host cell, stably transformed with a recombinant construct comprising the human FOXC2 promoter, in the presence and absence of a candidate agent, wherein an effect on the test level of expression as compared

20

25

30

1)

to control level of expression is indicative of an agent capable of regulating *FOXC2* promoter activity.

Methods for identification of polypeptides regulating FOXC2 promoter activity could include various techniques known in the art, such as the yeast one-hybrid system (see: Li & Herskowitz (1993) Science 262, 1870-1874) to identify proteins binding specific sequences from the FOXC2 regulatory region, biochemical purification of proteins which bind to the regulatory region, the use of a "southwestern" cloning strategy (see e.g. Hai et al. (1989) Genes & Development 3: 2083-2090) in which a pool of bacteria infected with a "phage library" are induced to express the encoded protein and probed with radioactive DNA sequences from the FOXC2 regulatory regions to identify binding proteins.

In a further aspect, the invention provides a human *FOXC2* enhancer region comprising a sequence selected from:

- (a) the nucleotide sequence set forth as positions 216 to 475, such as positions 223 to 231, positions 359 to 375, positions 378 to 402, or positions 403 to 423, in SEQ ID NO: 1, or a fragment thereof exhibiting FOXC2 enhancer activity;
  - (b) the complementary strand of (a); and
  - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).

As described above for the human FOXC2 promoter region, the invention further provides a recombinant construct comprising a human FOXC2 enhancer region, a vector comprising the said recombinant construct, as well as a host cell stably transformed with said vector or with said recombinant construct.

Further, the invention provides a method for identification of an agent regulating FOXC2 enhancer activity, said method comprising the steps: (i) contacting a candidate agent with the human FOXC2 enhancer region as defined above; and (ii) determining whether said candidate agent modulates expression of the FOXC2 gene, such modulation being indicative for an agent capable of regulating FOXC2 enhancer activity. It will be understood by the skilled person that known steps are available for performing such a method. For instance, a "panel" of constructs which include a variety of mutations and

deletions can be used in order to associate a response with a specific alteration of a single base or subsegment of the regulatory apparatus. A simple panel might include: enhancer plus promoter, promoter only, enhancer plus a "minimal" promoter from a distinct gene. As mentioned above, a transfection assay, using a host cell stably transformed with a suitable recombinant construct, can be a particularly useful screening assay for identifying an effective agent.

In yet a further aspect, the invention provides a method for identification of an agent capable of regulating a mammalian *FOXC2* promoter activity, said method comprising the steps (i) contacting a candidate agent with a murine *FoxC2* promoter nucleotide sequence shown as positions 216 to 2235, such as positions 216 to 475 or positions 1250 to 2235, in SEQ ID NO: 5; and (ii) determining whether said candidate agent modulates expression of a mammalian *FOXC2* gene, such modulation being indicative for an agent capable of regulating mammalian *FOXC2* promoter activity.

15

20

25

30

10

In another important aspect, the invention provides an isolated nucleic acid molecule selected from:

- (a) nucleic acid molecules comprising a nucleotide sequence as shown in SEQ ID NO: 3;
- (b) nucleic acid molecules comprising a nucleotide sequence capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence complementary the polypeptide coding region of a nucleic acid molecule as defined in (a) and which codes for a variant form of the FOXC2 transcription factor; and
- (c) nucleic acid molecules comprising a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence as defined in (a) or (b) and which codes for a variant form of the FOXC2 transcription factor.

In a preferred form of the invention, the said nucleic acid molecule has a nucleotide sequence identical with SEQ ID NO: 3 of the Sequence Listing. However, the nucleic acid molecule according to the invention is not to be limited strictly to the sequence shown as SEQ ID NO: 3. Rather the invention encompasses nucleic acid molecules carrying modifications like substitutions, small deletions, insertions or inversions, which nevertheless encode proteins having substantially the biochemical activity of the FOXC2 polypeptide according to the invention. Included in the invention are consequently nucleic

20

25

30

H

acid molecules, the nucleotide sequence of which is at least 90% homologous, preferably at least 95% homologous, with the nucleotide sequence shown as SEQ ID NO: 3 in the Sequence Listing.

Included in the invention is also a nucleic acid molecule which nucleotide sequence is degenerate, because of the genetic code, to the nucleotide sequence shown as SEQ ID NO: 3. A sequential grouping of three nucleotides, a "codon", codes for one amino acid. Since there are 64 possible codons, but only 20 natural amino acids, most amino acids are coded for by more than one codon. This natural "degeneracy", or "redundancy", of the genetic code is well known in the art. It will thus be appreciated that the nucleotide sequence shown in the Sequence Listing is only an example within a large but definite group of sequences which will encode the variant FOXC2 polypeptide.

The invention includes an isolated polypeptide encoded by the nucleic acid as defined above. In a preferred form, the said polypeptide has an amino acid sequence according to SEQ ID NO: 4 of the Sequence Listing. However, the polypeptide according to the invention is not to be limited strictly to a polypeptide with an amino acid sequence identical with SEQ ID NO: 4 in the Sequence Listing. Rather the invention encompasses polypeptides carrying modifications like substitutions, small deletions, insertions or inversions, which polypeptides nevertheless have substantially the biological activities of the variant FOXC2 polypeptide.

A further aspect of the invention is a vector harboring the nucleic acid molecule according to the invention. The said vector can e.g. be a replicable expression vector, which carries and is capable of mediating the expression of a DNA molecule according to the invention. In the present context the term "replicable" means that the vector is able to replicate in a given type of host cell into which is has been introduced. Examples of vectors are viruses such as bacteriophages, cosmids, plasmids and other recombination vectors. Nucleic acid molecules are inserted into vector genomes by methods well known in the art.

Included in the invention is also a cultured host cell harboring a vector according to the invention. Such a host cell can be a prokaryotic cell, a unicellular eukaryotic cell or a cell derived from a multicellular organism. The host cell can thus e.g. be a bacterial cell such as

10

15

20

30

11

an E. coli cell; a cell from yeast such as Saccharomyces cervisiae or Pichia pastoris, or a mammalian cell. The methods employed to effect introduction of the vector into the host cell are standard methods well known to a person familiar with recombinant DNA methods.

In yet another aspect, the invention includes a method for identifying an agent capable of regulating expression of the nucleic acid molecule as defined above, said method comprising the steps (i) contacting a candidate agent with the said nucleic acid molecule; and (ii) determining whether said candidate agent modulates expression of the said nucleic acid molecule.

In another aspect the invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to RNA transcribed by the alternatively spliced nucleic acid molecule shown as SEQ ID NO: 3, so as to prevent translation of the said RNA. Antisense nucleic acids (preferably 10 to 20 base-pair oligonucleotides) capable of specifically binding to control sequences for the alternatively spliced *FOXC2* gene are introduced into cells, e.g. by a viral vector or colloidal dispersion system such as a liposome. The antisense nucleic acid binds to the target nucleotide sequence in the cell and prevents transcription and/or translation of the target sequence. Phosphorothioate and methylphosphonate antisense oligonucleotides are specifically contemplated for therapeutic use by the invention. Suppression of expression of the alternatively spliced FOXC2 gene, at either the transcriptional or translational level, is useful to generate cellular or animal models for diseases/conditions related to lipid metabolism.

- In yet another aspect, the invention provides a method for the identification of polypeptides which bind to nucleotide sequences involved in the biological pathway regulating lipid metabolism and/or adipocyte differentiation, comprising the steps of:
  - (a) transfecting a host cell line with a human FOXC2 nucleotide sequence linked to a reporter gene, such as a gene encoding Green Fluorescent Protein (GFP) (for a review, see e.g. Galbraith et al. (1999) Methods in Cell Biology 58: 315-341);
  - (b) transfecting the said host cell line with a variety of human cDNA sequences, e.g. sequences included in a cDNA library;

- (c) identifying and isolating cells, e.g. by FACS cells sorting, having an altered level of expression of the said reporter gene, which is indicative that the polypeptide encoded by the added cDNA up- or downregulates at least one gene involved in the biological pathway regulating lipid metabolism and/or adipocyte differentiation;
- or a CRE-LOX mediated procedure (see e.g. Sauer (1998) Methods 14: 381-392); and (e) identifying the polypeptide expressed by the cDNA recovered in step (d), e.g. by sequencing the cDNA and comparing the obtained sequence against sequence databases.
- Throughout this description the terms "standard protocols" and "standard procedures", when used in the context of molecular biology techniques, are to be understood as protocols and procedures found in an ordinary laboratory manual such as: Current Protocols in Molecular Biology, editors F. Ausubel et al., John Wiley and Sons, Inc. 1994, or Sambrook, J., Fritsch, E.F. and Maniatis, T., Molecular Cloning: A laboratory manual,

  2<sup>nd</sup> Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY 1989.

## **EXAMPLES**

Additional features of the invention will be apparent from the following Examples.

Examples 1 to 4 are actual, while the remaining Examples are prophetic.

# EXAMPLE 1: Computational identification of FOXC2 genomic sequences

The sequences present in the GenBank database (http://www.ncbi.nlm.nih.gov) were screened for sequence similarity to the human FOXC2 cDNA sequence (GenBank accession number NM\_00521 (SEQ ID NO: 9)). The BLAST algorithm (Altschul et al. (1997) Nucleic Acids Res. 25:3389-3402) was used for determining sequence identity. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov). A working draft genomic sequence in 25 unordered pieces, from the Homo sapiens chromosome 16 clone RP11-463O9 (GenBank accession number AC009108; Version 6; GI:7689930; released 4 May 2000), was selected for further studies.

Regions in sequence AC009108 matching portions of the FOXC2 cDNA sequence NM\_005251were combined using the PHRAP software, developed at the University of Washington (http://www.genome.washington.edu/UWGC/analysistools/phrap.htm). Two contigs of 9780 bp (positions 116445 to 126224 in GenBank AC009108.6) and 3784 bp (positions 42927 to 46710 in GenBank AC0091108.6), respectively, were assembled to generate a human FOXC2 genomic fragment of 13451 bp.

The ClustalW multiple sequence alignment program, version 1.8 (Thompson et al. (1994) Nucleic Acids Research 22: 4673-4680), was then used to identify the human *FOXC2* extended genomic DNA sequence of 6458 bp (SEQ ID NO: 1) by comparison with the mouse cDNA sequence X74040 (SEQ ID NO: 6). First, a 6459 bp sequence, corresponding to positions 1500-7958 in the 13451 bp sequence, was selected. Positions 1-2285 in this 6459 bp sequence corresponded to 44426-46710 in AC009108.6, while positions 2151-6459 corresponded to positions 126224-121916 (reverse complement taken) in AC009108.6. The overlap of positions 2151-2285 allowed for the contigs to be joined by the assembly program. The G residue in position 2655 was considered to be a sequencing error and was removed, which resulted in the 6458 bp sequence set forth as SEQ ID NO: 1. The open reading frame in SEQ ID NO: 1 encodes a polypeptide (SEQ ID NO: 2) identical with the known human FOXC2 polypeptide shown as SEQ ID NO:10.

EXAMPLE 2: Identification of potential regulatory sequences in the human and mouse *FOXC2* genomic sequences

In phylogenetic footprinting (for a review, see Duret & Bucher (1997) Current Opinion in Structural Biology 7(3): 399-406) sequences are aligned and a regional sequence identity is determined for each window of a fixed, arbitrary length. This allows the identification of potential regulatory regions in genomic sequences. Non-exon sequences that are conserved over the course of evolution are likely to perform regulatory roles. Phylogenetic footprinting was performed as described in Wasserman & Fickett (1998) J. Mol. Biol. 278, 167-181, based on an alignment generated with the ClustalW multiple sequence alignment program, version 1.8 (Thompson et al. (1994) Nucleic Acids Research 22: 4673-4680),

25

20

10

15

20

25

30

with default parameters adjusted to a gap opening penalty of 20 and a gap extension penalty of 0.2. The human (SEQ ID NO: 1) and mouse (SEQ ID NO: 5) genomic sequences were aligned. Percentage identity was plotted for each contiguous 200 bp segment of the human gene to identify segments differentially conserved (in comparison to adjoining sequences) (Fig. 2).

In addition to segments of the published exon sequence, two differentially conserved regions or "footprints" were identified in the human gene. Both of these regions are local maxima and contain segments which exceed 70% nucleotide identity between the human and mouse genomic sequences. One region, shown as positions 1250 to 2235, in particular positions 1250 to 1749, in SEQ ID NO: 1, immediately adjacent to the published exon region, is likely to contain the transcription start site and proximal promoter regulatory sequences (Fig. 4). Another region, shown as positions 216 to 475 in SEQ ID NO: 1, approximately 1700 bp distal from the transcription start site, is likely to function as some form of regulatory region (either enhancer or repressor) (Fig. 3). (A schematic overview of the extended *FOXC2* gene is shown in Fig. 1.)

Further analysis of these regulatory regions identified short segments of higher conservation between the mouse and human genes, suggesting that these specific segments function as transcription factor binding sites. The *TRANSFAC* transcription factor database (http://transfac.gbf.de) (see Wingender et al. (2000) Nucleic Acids Research 28(1): 316-319) was screened for matches to known transcription factors. Consensus sites (identifiers R05066; R05067; R05068; and R05069) were found to match sequences conserved between the human *FOXC2* and mouse FoxC2 genes. This suggests the presence of multiple forkhead-like binding sites in the distal regulatory enhancer, and potential autoregulation of FOXC2 by its protein product.

The same analysis was performed with reference to 200 bp contiguous segments of the mouse *FoxC2* genomic sequence (SEQ ID NO: 5). The following conserved regions were identified: 190 to 420; 1070 to 1645; and 5580 to 5875. They correlate to the regions indicated above for the human sequence and should be considered orthologous regions.

15

20

25

## EXAMPLE 3: Identification of an alternative human FOXC2 cDNA sequence

BLASTN screening of the dbEST database from GenBank, using the human FOXC2 cDNA (SEQ ID NO: 9) as a query sequence, revealed several ESTs overlapping containing portions of the available cDNA. A specialized tool, est\_genome (http://www.sanger.ac.uk), for the prediction of exon boundaries using ESTs was applied to compare the EST sequences to the genomic sequences (See Mott, R. (1997) Computer Applications in the Biosciences 13(4): 477-478). Two classes of ESTs were observed: sequences extending into the 3'-untranslated region and sequences revealing an alternative first exon spliced to a junction internal to the previously described first exon.

Specifically, it was found that the nucleotides in positions 33 to 182 in the EST with accession no. AW271272 (SEQ ID NO: 11) were identical to positions 66 to 215 in the extended *FOXC2* genomic sequence (SEQ ID NO: 1), and that positions 183 to 327 in SEQ ID NO: 11 were identical to positions 2516 to 2660 in SEQ ID NO: 1. Similarly, positions 5 to 55 in the EST with accession no. AW793237 (SEQ ID NO: 12) were identical to positions 165 to 215 in the extended *FOXC2* genomic sequence (SEQ ID NO: 1), and positions 56 to 157 in SEQ ID NO: 12 were identical to positions 2516 to 2607 in SEQ ID NO: 1. These results revealed an alternative splicing pattern in the human *FOXC2* gene. According to this splicing pattern, an alternative gene sequence (SEQ ID NO: 3) is derived by joining the regions shown as positions 1–215 and 2516–6458 in SEQ ID NO: 1. Alternative splicing patterns are known to regulate the synthesis of a variety of peptides and proteins. It may result in proteins with an entirely different function or in dysfunctional or inhibitory splice products (for a review, see McKeown (1992) Annu. Rev. Cell. Biol. 8: 133-155).

The amino acids corresponding to positions 1 to 94 in the published FOXC2 transcription factor (SEQ ID NO: 10) are missing in protein encoded by the spliced variant generated from the alternative promoter (SEQ ID NO: 4). Consequently, the entire region N-terminal of the DNA binding domain and a portion of the DNA-binding domain (corresponding to positions 72-94 in SEQ ID NO: 2) are not present in the splice variant. It is postulated that this truncation leads to a protein which has a deficient "forkhead" DNA-binding region, and thus has a potential inhibitory function on the biological activities of the FOXC2

15

20

sequence.

protein. This truncated FOXC2 protein may have a role in regulation of FOXC2, and an involvment in adipocyte differentiation and adipogenesis.

## EXAMPLE 4: Cloning and sequencing of the FOXC2 promoter

The DNA region corresponding to nucleotide 176 to nucleotide 2233 (SEQ ID NO. 1 version 2) has been cloned using nested PCR on human genomic DNA. The PCR was performed according the Herculase™ protocol (Stratagene catalog #600260; http://www.stratagene.com/pcr/herculase.htm) and with the inclusion of 8-10% DMSO.

In the initial reaction, the 5'-primer KRKX131 (CCATTGCCTTCTAGTCGCCTCC) was used together with the 3'-primer KRKX133 (CGTTGGGGTCGGACACGGAGTA) using 250 ng Clontech Genomic DNA # 6550-1 as template. The nested reaction was performed on 1/100 of the initial PCR reaction using the 5'-primer KRKX132 (GGTACCTACGCAGCCGATGAACAGCCA) and the 3'-primer KRKX134 (GCTAGCGCTGCTTCCGAGACGGCTCG). After the second PCR, the product was analyzed by electrophoresis in a 1.2% agarose gel, and a PCR product of the expected size was obtained and extracted for ligation into a TOPO PCR2.1 vector (Invitrogen, Carlsbad, CA) by standard cloning procedures and thereafter sequenced. The PCR reaction and cloning procedure was repeated in two parallel separate experiments, and sequence data from the two separate reactions were compared with the bioinformatically assembled

A DNA region containing the promoter (Fig. 4) corresponding to nt1179 to 2233 (SEQ ID NO: 1, version 2) was has been cloned using nested PCR in the same manner as described above. In the initial reaction, the 5'-primer KRKX136 (GGTACCCCCGAGCCTGGAAACTCCCT) was used together with the 3'-primer KRKX134 (GCTAGCGCTGCTTCCGAGACGGCTCG) using 250 ng genomic DNA as a template. The PCR reaction and cloning procedure was repeated in four parallel separate experiments, and sequence data from the four separate reactions were compared with the bioinformatically assembled sequence.

# EXAMPLE 5: Tissue expression profiling of the alternative transcript

Tissue expression profiling of the alternative transcript (SEQ ID NO: 3) is performed using standard Northern blotting procedures. RNA samples from an array of human tissues, including adipose tissue, are analyzed using an RNA or DNA probe specific for the alternative transcript. The expression profile in adipose tissue could be indicative a putative regulatory function for the alternative gene product (SEQ ID NO: 4) on adipogenesis and adipocyte differentiation.

10

5

In addition, reverse transcriptase PCR (RT-PCR) according to standard procedures is used to detect very low level expression of the alternative transcript in adipose tissue. RNA is prepared from human adipose tissue, and RT-PCR is performed using PCR primers specific for the alternative transcript.

15

## EXAMPLE 6: Mapping of the 5'-edge of the alternative exon by RACE-PCR

RNA is prepared from human adipose tissue using standard protocols. RACE (Rapid
Amplification of cDNA Ends) PCR is performed using the SMART<sup>TM</sup> RACE cDNA
Amplification Kit (Clontech catalogue No. K1811-1; http://www.clontech.com/product/
catalog/PCR/smartrace.html). With this procedure, the first strand synthesis produces
cDNA with an extension containing a known sequence. Due to the mechanism of the
extension procedure, the extension is typically added only to complete first strand cDNAs.

The 5'-RACE PCR is then performed using the 5'-primer provided with the kit, together with a 3'-primer corresponding to positions 210-237 in SEQ ID NO: 3

(GAACTGGTAGATGCCGTTCAAGGTTTCC) specific for the alternative transcript. The PCR product is cloned into a cloning vector and sequenced using standard protocols.

25

30

### **EXAMPLE 7: Functional analysis**

The identified regulatory regions are analyzed to determine their impact on the transcription of the FOXC2 gene or a reporter gene substituted for FOXC2. A PCR reaction is performed to isolate the promoter region adjacent to the published exon sequence, possibly including the sequences extending to the beginning of the ATG encoding the first methionine. This PCR product is cloned into a reporter plasmid adjacent to a reporter gene (e.g. luciferase). The upstream regulatory region, i.e. regions containing both upstream and promoter proximal sequences, or these sequences bearing artificially induced differences, are cloned in a similar manner. These constructs are transfected into a cell culture model system and the level/activity of the protein encoded by the reporter gene is determined. This would provide information on the function of the identified regions, and used to assess the impact of the different regions on transcriptional regulation. Similarly, the upstream regulatory region, a region containing both upstream and promoter proximal sequences, or these sequences bearing artificially induced differences can be cloned and used to assess the impact of these regions on the transcription of the reporter gene.

### 20 EXAMPLE 8: Reporter gene assay to identify modulating compounds

Reporter gene assays are well known as tools to signal transcriptional activity in cells. (For a review of chemiluminescent and bioluminescent reporter gene assays, see Bronstein et al. (1994) Analytical Biochemistry 219, 169-181.) For instance, the photoprotein luciferase provides a useful tool for assaying for modulators of promoter activity. Cells are transiently transfected with a reporter construct which includes a gene for the luciferase protein downstream from the FOXC2 promoter and enhancer region, or fragments thereof regulating the FOXC2 activity. Luciferase activity may be quantitatively measured using e.g. luciferase assay reagents that are commercially available from Promega (Madison, WI). Differences in luminescence in the presence versus the absence of a candidate modulator compound are indicative of modulatory activity.

TABLE I
Summary of FOXC2 sequences

SEQ ID NO:	GenBank accession no.	Description
1		Human FOXC2 extended genomic DNA sequence
2		Human FOXC2 polypeptide sequence (Identical with SEQ ID NO: 10)
3		Human FOXC2 DNA sequence Alternative splicing
4		Human polypeptide sequence Alternative open reading frame
5	Y08222	Mouse MHF-1 (FaxC2) genomic DNA sequence (CDS 2070 – 3554)
6	X74040	Mouse MHF-1 (FoxC2) cDNA sequence
7		Mouse MHF-1 (FaxC2) polypeptide sequence
8	Y08223	Human FKHL14 (FOXC2) genomic DNA sequence (CDS 1197 – 2702)
9	NM_005251	Human FKHL14 (FOXC2) cDNA sequence
10		Human FKHL14 (FOXC2) polypeptide sequence
11	AW 271272	Human EST
12	AW 793237	Human EST




TABLE II
Summary of features in human FOXC2 sequences shown as SEQ ID NOs: 1 and 3

Feature	Positions
SEQ ID NO: 1	
First exon according to the alternative transcript	1-215
- Untranslated region	1 – 186
- Region coding for 5'-part of alternative protein	187 – 215
Alternative first exon splice site	215 – 216
Predicted enhancer region	216 – 475
– E-box-like region	223 – 231
– Forkhead-like region	359 – 375
– Forkhead-like region	378 – 402
– Ets-like region	403 – 423
Predicted promoter region	1250 – 1749
– Forkhead-like region	1692 - 1703
First exon according to the published form of the transcript	1746 – 4629
- Untranslated region	1746 – 2234
- Polypeptide coding region	2235 – 3740
- Region coding for DNA-binding domain	2448 – 2735
Second exon according to the alternative transcript	2516 – 4629
- Portion of polypeptide used in alternative transcript	2516 - 3740
- Untranslated region	3741 - 4629
SEQ ID NO: 3	
Polypeptide coding region (5' of splice site)	187 – 215
Polypeptide coding region (3' of splice site)	216 – 1437
- Region coding for truncated portion of protein	216 - 435



#### **CLAIMS**

5

25

- 1. A human FOXC2 promoter region comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 1692 to 1703 in SEQ ID
- NO: 1, or a fragment thereof exhibiting FOXC2 promoter activity;
  - (b) the complementary strand of (a); and
  - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
- The human *FOXC2* promoter region according to claim 1, comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 1250 to 1749 in SEQ ID
  - NO: 1, or a fragment thereof exhibiting FOXC2 promoter activity;
  - (b) the complementary strand of (a); and
- (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
  - 3. The human FOXC2 promoter region according to claim 2, comprising a sequence selected from:
- (a) the nucleotide sequence set forth as positions 1250 to 2235 in SEQ ID
   NO: 1, or a fragment thereof exhibiting FOXC2 promoter activity;
  - (b) the complementary strand of (a); and
  - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
  - A recombinant construct comprising the human FOXC2 promoter region according to any one of claims 1 to 3.
  - 5. The recombinant construct according to claim 4 wherein the human *FOXC2* promoter region is operably linked to a gene encoding a detectable product.
  - The recombinant construct according to claim 5 wherein said gene encoding a
    detectable product is a human FOXC2 gene.

15

20

- 7. The recombinant construct according to claim 4 further comprising a reporter gene.
- 8. A vector comprising the recombinant construct according to any one of claims 4 to 7.
- A host cell stably transformed with the recombinant construct according to any one of claims 4 to 7.
- 10. A method for identification of an agent regulating *FOXC2* promoter activity, said method comprising the steps
  - (i) contacting a candidate agent with a human FOXC2 promoter region as defined in any one of claims 1 to 3; and
  - (ii) determining whether said candidate agent modulates expression of the FOXC2 gene, such modulation being indicative for an agent capable of regulating FOXC2 promoter activity.
  - 11. A method for identification of an agent regulating FOXC2 promoter activity, said method comprising assaying reporter gene expression in a cell according to claim 9 in the presence and absence of a candidate agent, wherein an effect on the test level of expression as compared to control level of expression is indicative of an agent capable of regulating FOXC2 promoter activity.
  - 12. A human FOXC2 enhancer region comprising a sequence selected from:
    - (a) the nucleotide sequence set forth as positions 223 to 231 in SEQ ID NO: 1, or a fragment thereof exhibiting FOXC2 enhancer activity;
    - (b) the complementary strand of (a); and
    - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
- 30 13. A human FOXC2 enhancer region comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 359 to 375 in SEQ ID NO: 1, or a fragment thereof exhibiting FOXC2 enhancer activity;
  - (b) the complementary strand of (a); and

- (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
- 14. A human FOXC2 enhancer region comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 378 to 402 in SEQ ID NO: 1, or a fragment thereof exhibiting FOXC2 enhancer activity;
    - (b) the complementary strand of (a); and
    - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).

5

- 15. A human FOXC2 enhancer region comprising a sequence selected from:
  - (a) the nucleotide sequence set forth as positions 403 to 423 in SEQ ID NO: 1, or a fragment thereof exhibiting *FOXC2* enhancer activity;
  - (b) the complementary strand of (a); and
- (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).
  - 16. The human *FOXC2* enhancer region according to any one of claims 12 to 15 comprising a sequence selected from:
    - (a) the nucleotide sequence set forth as positions 216 to 475 in SEQ ID NO: 1, or a fragment thereof exhibiting FOXC2 enhancer activity;
      - (b) the complementary strand of (a); and
      - (c) nucleotide sequences capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence as defined in (a) or (b).

25

20

- 17. A recombinant construct comprising a human *FOXC2* enhancer region according to any one of claims 12 to 15.
- 18. A vector comprising the recombinant construct according to claim 17.

30

19. A host cell stably transformed with the recombinant construct according to claim 18.

20

25

- A method for identification of an agent regulating FOXC2 enhancer activity, said method comprising the steps
  - (i) contacting a candidate agent with the human FOXC2 enhancer region as defined in any one of claims 12 to 16; and
- (ii) determining whether said candidate agent modulates expression of the *FOXC2* gene, such modulation being indicative for an agent capable of regulating *FOXC2* enhancer activity.
- 21. A method for identification of an agent capable of regulating FOXC2 enhancer
  activity, said method comprising assaying reporter gene expression in a cell as
  defined in claim 19 in the presence and absence of a candidate agent, wherein an
  effect on the test level of expression as compared to control level of expression is
  indicative of an agent capable of regulating FOXC2 enhancer activity.
- 15 22. A method for identification of an agent capable of regulating a mammalian *FOXC2* promoter activity, said method comprising the steps
  - (i) contacting a candidate agent with a murine FoxC2 promoter nucleotide sequence shown as positions 1250 to 2235 in SEQ ID NO: 5; and
  - (ii) determining whether said candidate agent modulates expression of a mammalian *FOXC2* gene, such modulation being indicative for an agent capable of regulating mammalian *FOXC2* promoter activity.
    - 23. A method for identification of an agent capable of regulating a mammalian *FOXC2* enhancer activity, said method comprising the steps
      - (i) contacting a candidate agent with a murine FoxC2 enhancer nucleotide sequence shown as positions 216 to 475 in SEQ ID NO: 5; and
      - (ii) determining whether said candidate agent modulates expression of a mammalian *FOXC2* gene, such modulation being indicative for an agent capable of regulating mammalian *FOXC2* enhancer activity.
    - 24. A method for identification of an agent capable of regulating a mammalian *FOXC2* enhancer activity, said method comprising the steps

- (i) contacting a candidate agent with a murine FoxC2 enhancer nucleotide sequence shown as positions 216 to 2235 in SEQ ID NO: 5; and
- (ii) determining whether said candidate agent modulates expression of a mammalian *FOXC2* gene, such modulation being indicative for an agent capable of regulating mammalian *FOXC2* enhancer activity.
- 25. An isolated nucleic acid molecule selected from:
  - (a) nucleic acid molecules comprising a nucleotide sequence as shown in SEQ ID NO: 3;
- (b) nucleic acid molecules comprising a nucleotide sequence capable of hybridizing, under stringent hybridization conditions, to a nucleotide sequence complementary the polypeptide coding region of a nucleic acid molecule as defined in (a) and which codes for a variant form of the FOXC2 transcription factor; and (c) nucleic acid molecules comprising a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence as defined in (a) or (b) and
  - 26. An isolated polypeptide encoded by the nucleic acid according to claim 25.

which codes for a variant form of the FOXC2 transcription factor.

- 27. The isolated polypeptide according to claim 26 having an amino acid sequence shown as SEQ ID NO: 4 in the Sequence Listing
  - 28. A vector harboring the nucleic acid molecule according to claim 25.
- 25. A replicable expression vector, which carries and is capable of mediating the expression of a nucleotide sequence according to claim 25.
  - 30. A cultured host cell harboring a vector according to claim 28 or 29.
- 31. A process for production of a variant form of the FOXC2 transcription factor polypeptide, comprising culturing a host cell according to claim 30 under conditions whereby said polypeptide is produced, and recovering said polypeptide.

10

- 32. A method for identifying an agent capable of regulating expression of the nucleic acid molecule according to claim 25, said method comprising the steps
  - (i) contacting a candidate agent with the said nucleic acid molecule; and
  - (ii) determining whether said candidate agent modulates expression of the said nucleic acid molecule.
- 33. An antisense oligonucleotide having a sequence capable of specifically hybridizing to RNA transcribed by the nucleic acid molecule according to claim 25, so as to prevent translation of the said RNA.
- 34. A method for the identification of polypeptides which bind to nucleotide sequences involved in the biological pathway regulating lipid metabolism and/or adipocyte differentiation, comprising
  - (a) transfecting a host cell line with a human FOXC2 nucleotide sequence linked to a reporter gene;
  - (b) transfecting the said host cell line with a variety of human cDNA sequences;
  - (c) identifying and isolating cells having an altered level of expression of the said reporter gene;
  - (d) recovering cDNA from the cells isolated in step (c); and
- 20 (e) identifying the polypeptide expressed by the cDNA recovered in step (d).

### **ABSTRACT**

The present invention relates an isolated promoter region of the mammalian transcription factor FOXC2. The invention also relates to screening methods for agents modulating the expression of FOXC2 and thereby being potentially useful for the treatment of medical conditions related to obesity. The invention further relates to a previously unknown variant of the human FOXC2 gene, derived via the use of an alternative promoter, which produces an additional exon that generates a distinct open reading frame via splicing. The alternative gene encodes a variant of the FOXC2 transcription factor, which is lacking a part of the DNA-binding domain and consequently has a potential regulatory function.

### SEQUENCE LISTING

<110> Pharmacia & Upjohn AB <120> Promoter Sequences <130> 00298 <140> <141> <160> 12 <170> PatentIn Ver. 2.1 <210> 1 <211> 6458 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (2235)..(3740) cctttggctt tgaattgatc aggagacaaa gataatgcat ctacattttc gtcttctgtt 60 cttttattgq aaataagtgg cacgcccat tgccttctag tcgcctcccc gaagcgaaga 120 qqccqaaqcq aaqaqqcctg gtqggttgtc tcaacatcct tttgctgaga atcgaatacg 180 cagoogatga acagocagga agggtgcaag gaaacotgaa atacaaatgt totocotgaa 240 geoetettee etgeecaace agaceageaa ettecaaaat tetgeeegtg tttageettg 300 ttaaagqqqt gtctcactcc ttcagqqaaa gtqgqaaaag gqqatctgat tattqaqqtq 360 tqqaaqqaat aaataatcag tccacaaata aacaaactgt ccgggattcc tagagggaag 420 qaqaaatcct tqaaqqaqat ccaaqtcqct ccaqqtctqc ctqccqaata atatcatccc 480 gaagggatet tgaaccgttt gcaatcaacc geteacceag tetteccaeg gagegegete 540 cctaactcac cctacccacc caacaaaaca aaaaaaaggc tgaaatatag aaaagcaact 600 tgqaggctcc cagggggacg ttgccaggag caggaggcag ggacagcgcc ctagggtcgg 660 tgttagegge eggeeegge etgggeeaeg ggaaaegtee aegettggtg eeegeggtge 720 geggegetea ttgegegege ettegageea ageeeeegeg gaaaaeagge tegggtttet 780 cctcgcaggg cccaggaact cggctctgcc tggcccgggt gggtcgctgc attgtcccgg 840 tettetggga gtgeggggte agettgttag agggaattte taeetgggaa aagggagaeg 900 agtttcgaag ctgaagttgg taggctgcga gtgtccacgc gggagacgaa agggggaaat 960

agcagagtca cttcaccett ttececaaac eccacaaaac tgetegcage gaegeggatg 1020

.1

atctaccqaa ttccccqcqa attcqqaqqa ttaagttqtc agtcagcacg ttgctacctt 1080 cccctctatg cactccgctg cctggctcct cggcggggag cgagggaaac tcagtttgta 1140 gggtttacct ctaaaacctc gataggttat ccttgacgac cccgagcctg gaaactccct 1200 gttgatgatt aattatttga ttaaataagt ataacatcca ggagaggccc tgccattcca 1260 atccagegeg tttgettttg aatccattae acctgggeee ceataattag gaaatetaat 1320 tattcgcttc atcactcatt aataagaaaa atgtcccagg atcattgcta cttacaaggt 1380 ctttgggaga gatattttac tctattaatc cattctattt tatatttcaa attgattttt 1440 tttaacagag gaaaqtggct atctttttgt tttgggcatg tgggcccatt caccaaaatg 1500 tgatcataaa ataaatttta ataagatata actttttaaa aagttttcaa gtgaagacgg 1560 agtogoogeg gaggoogggg oggoggggto ttagagooga oggattootg ogctootogo 1620 cccgattggc gccggactcc tctcagctgc cgggtgattg gctcaaagtt ccgggagggg 1680 gcgtggcccg aggaaagtaa aaactcgctt tcagcaagaa gacttttgaa acttttccca 1740 atocotaaaa gggacttggo ototttttot gggotcagog gggcagoogo toggaccoog 1800 gegegetgae cetegggget geegattege tgggggettg gagageetee tgegeeeete 1860 ctcgcgcggg ccgaqggtcc accttggtcc ccaggccgcg gcgtctccgc tgggtccgcg 1920 geogeoegee tgeoegeget geogeogeeg ggteetggag ceagegagga geggggeegg 1980 cgctgcgctt gcccggggcg cgccctccag gatgccgatc cgcccggtcc gctgaaagcg 2040 cgcgcccctg ctcggcccga gcgacgacga ccgcgcaccc tcgccccgga ggctgccagg 2100 agaccggggc cgcccctccc gctcccctcc tctccccctc tggctctctc gcgctctctc 2160 qctctcaggq ccccctcgc tcccccqqcc qcagtccgtg cgcqaqggcg ccgqcqagcc 2220 gtotoggaag cago atg cag gog ogo tao too gtg too gao oco aac goo Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala ctg gga gtg gtg ccc tac ctg agc gag cag aat tac tac cgg gct gcg 2318 Leu Gly Val Val Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala 15 gge age tac gge gge atg gee age eee atg gge gte tat tee gge eae 2366 Gly Ser Tyr Gly Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His ccg gag cag tac agc gcg ggg atg ggc cgc tcc tac gcg ccc tac cac 2414 Pro Glu Gln Tyr Ser Ala Gly Met Gly Arg Ser Tyr Ala Pro Tyr His 2462 cac cac cag ccc gcg gcg cct aag gac ctg gtg aag ccg ccc tac agc His His Gln Pro Ala Ala Pro Lys Asp Leu Val Lys Pro Pro Tyr Ser 70 75 65

. i

	atc Ile			Ile		-	_		_					_	_	2510
	acc Thr															2558
	cgg Arg 110			_	_			_		-		-				2606
	ctc Leu															2654
	aag Lys		_				-		-	-				_		2702
	aac Asn		_		_		_			_			_	_	-	2750
	tcc Ser	_		-				-			_		_		_	2798
	gcg Ala 190		_		_	_	-						-	_		2846
	gag Glu															2894
	ctg Leu	_	_			-			_	_	_		-	_		2942
_	cag Gln		_	_	_	_		-		-		_				2990
	ggt Gly															3038
	ttc Phe 270															3086
	gag Glu															3134

	gcg Ala															3182
	gct Ala															3230
	cga Arg															3278
tgc Cys	gtc Val 350	ccg Pro	ccc Pro	gcc Ala	ctg Leu	gac Asp 355	gag Glu	gcc Ala	ctc Leu	tcg Ser	gac Asp 360	cac His	ccg Pro	agc Ser	ggc Gly	3326
	acg Thr															3374
	ctc Leu															3422
	cag Gln															3470
	ccc Pro															3518
	ggg Gly 430															3566
	act Thr				-			_							–	3614
	gag Glu															3662
	cag Gln															3710
	tac Tyr			_	-	-			tga	cgtg	jteco	gg g	jacct	cccc	:t	3760
ccc	egged	cg c	tccç	gctt	c go	ttcc	cago	ccc	gaco	caa	ccaç	jacaa	itt a	aggg	gctgc	3820
aga	gacgo	caa a	aaaq	gaaac	a aa	acat	gtco	acc	aacc	ttt	tete	agac	ေင င	ggaç	cagag	3880
agc	gggca	acg c	tago	cccc	a go	egte	etgtg	, aag	agco	cag	gtaa	cttt	aa t	tcgc	cgccc	3940
cgt	ttctc	igg a	tcc	agga	a ac	ccct	ccaa	agg	gaco	cag	ccca	acaa	aa t	gagt	attgg	4000

::

tettaaaate ecceteeet accaggaegg etgtgetgtg etegacetga gettteaaaa 4060 gttaagttat ggacccaaat cccatagega gcccctagtg actttctgta ggggtcccca 4120 taggtgtatg ggggtctcta tagataatat atgtgctgtg tgtaatttta aatttctcca 4180 acceptgctgt acaaatgtgt ggatttgtaa tcaggctatt ttgttgttgt tgttgtttgtt 4240 cagagocatt aatataatat ttaaagttga gttcactgga taagtttttc atcttgccca 4300 accatttcta actqccaaat tgaattcaaq aaaccgatqt qqqttttqtt tcctqtacaa 4360 ttatqaqata taattetttt teecattgta ggtettttae aaaacaagaa aataatttat 4420 ttttttqttq qtqqataaag aagtcaagta tctqatactt tttatttaca aagtgtqatq 4480 gttttgtata gtaggttcca ccctgagtat tcctaaaaga aaaaaaaaa aaaagcttaa 4540 aaactctaac ttcatctgtg tttgtcttac gtggtcttaa tcgttgtact taccttaaaa 4600 taaacccatg ttgttttttc tgcccaaagt ttggacagtg tgtttgtgtt gttgcatttt 4660 ttacaaacga ggtgtgtttg caaacccacc tgctttgatt atttttgtta cacaggtggg 4720 tatatgtgta gacacataaa aacgaccaga gaataggagc acacacctgc tgtcttgttt 4780 agtgacagaa aaaggctttt gattaatttt aaaatcccac tetaggattt tttcttttcg 4840 agaaaccgcc cagttggagg gggctgcctg aaggaccgga ccatgagttt gccgtgatgc 4900 attiticttaa atgcacaaaa acatgctaat tgtcaaaaca aacagtgcca ctccatctca 4960 gtgtccagcc gtccccagtt taggaggtga aggaagggaa gaataaacat ttcccgtttg 5020 ctaactgcaa cccagggtga gtcctgcttt cccccgattt tataaaattt gagcctcttt 5080 gcctgcttta atagttttcc agagaatttg aactgggcca atgaaggtct gaaggggacg 5140 gattttctag cgtttgatat ccatcccct tagcggccag atcagagggg aatttcagac 5200 tttattactt ctcaatgtca tgtctaaatc tacaccetca tcgcagtgaa aaattttaaa 5260 acctcattac ccttcaaaaa taatttatga tatttttaga gttctaaatt caagtttttc 5320 aatatgttaa ataatagaga ttattttttg ttttcaatgt taatatctcg tcttttacat 5380 ttttaatagt aacatagttt ttgtgaaatg tagctgacga aatggcttta ttatctattt 5440 caatggctga agtccaccac teceetgetg geetetatgt gtgaatttgg ggaccaaage 5500 ttcatcaatt cccaccccag caggtgagct gtaccttgct aatgctgaag ttctttgtga 5560 gcttaacqtt tcaagaccag atgattttgc taaaggtgat tttgcttgat gcagtggcgc 5620 tgaacgtaac ccgggtgttt ttgtcgtgtt gttttcaaca tggcacttta tctccacgct 5680 atgttgaaat agaattaggg gaagcttaaa gcataataat tgtccccaca tgtgcaacac 5740

..

agactettic aatetytgge eccagagyty geacacaytt aagacttyge gyctytetea 5800 ttettttea taatytgegy gtteeegygt gteegygtye tagacttea geagyeecea 5860 gyccayacyg getttygtty agtgaacayg agyagyaayt taagyagyta gygytygyga 5920 gagaceetet ecaayetyca gaayaayyty geecaayete ettgeetyeg tetgeegya 5980 tyytteatt ttaettetye tegetteaty etattyeece cagyayaaya gygyagyattt 6040 ecayacygta agegagytyg etttteett teeetagaey tttaaaayaa atettetya 6100 aagetyeee teategtaay ettgaaace gttygtytee tyttagtyge gagygetyay 6160 agacacygy agaaataaay gagagegaey gtgtygetya gageeeccay gtetgetyt 6220 gaaactaaye tygyettity cacettay aageetttt aaayaaytee tyetytytyg 6280 gygeegyaay eccaaytyay tygyeetty gyagyttate gygagyyte tttaceacte 6340 ettgyggaac gtgygeaacy gygygattyt atetgaaget ttatteayy gacteate 6458

<210> 2 <211> 501 <212> PRT <213> Homo sapiens

210

<400> 2 Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala Leu Gly Val Val 10 Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala Gly Ser Tyr Gly 20 Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His Pro Glu Gln Tyr 35 40 45 Ser Ala Gly Met Gly Arg Ser Tyr Ala Pro Tyr His His His Gln Pro 55 Ala Ala Pro Lys Asp Leu Val Lys Pro Pro Tyr Ser Tyr Ile Ala Leu 65 70 Ile Thr Met Ala Ile Gln Asn Ala Pro Glu Lys Lys Ile Thr Leu Asn 90 85 Gly Ile Tyr Gln Phe Ile Met Asp Arg Phe Pro Phe Tyr Arg Glu Asn 100 105 110 Lys Gln Gly Trp Gln Asn Ser Ile Arg His Asn Leu Ser Leu Asn Glu 115 120 125 Cys Phe Val Lys Val Pro Arg Asp Asp Lys Lys Pro Gly Lys Gly Ser 130 135 140 Tyr Trp Thr Leu Asp Pro Asp Ser Tyr Asn Met Phe Glu Asn Gly Ser 150 155 Phe Leu Arg Arg Arg Arg Phe Lys Lys Asp Val Ser Lys Glu 165 170 Lys Glu Glu Arg Ala His Leu Lys Glu Pro Pro Pro Ala Ala Ser Lys 180 185 190 Gly Ala Pro Ala Thr Pro His Leu Ala Asp Ala Pro Lys Glu Ala Glu 195 200 205 Lys Lys Val Val Ile Lys Ser Glu Ala Ala Ser Pro Ala Leu Pro Val

Ile Thr Lys Val Glu Thr Leu Ser Pro Glu Ser Ala Leu Gln Gly Ser 235 230 Pro Arq Ser Ala Ala Ser Thr Pro Ala Gly Ser Pro Asp Gly Ser Leu 250 255 245 Pro Glu His His Ala Ala Ala Pro Asn Gly Leu Pro Gly Phe Ser Val 265 270 260 Glu Asn Ile Met Thr Leu Arg Thr Ser Pro Pro Gly Gly Glu Leu Ser 280 275 285 Pro Gly Ala Gly Arg Ala Gly Leu Val Val Pro Pro Leu Ala Leu Pro 295 300 Tyr Ala Ala Ala Pro Pro Ala Ala Tyr Gly Gln Pro Cys Ala Gln Gly 305 310 315 Leu Glu Ala Gly Ala Ala Gly Gly Tyr Gln Cys Ser Met Arg Ala Met 335 325 330 Ser Leu Tyr Thr Gly Ala Glu Arg Pro Ala His Met Cys Val Pro Pro 340 345 350 Ala Leu Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Thr Ser Pro 355 360 365 Leu Ser Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Ala Ala 370 375 380 Thr Gly His His Gln His His Gly His His Fro Gln Ala Pro 390 395 Pro Pro Pro Pro Ala Pro Gln Pro Gln Pro Thr Pro Gln Pro Gly Ala 405 410 415 Ala Ala Ala Gln Ala Ala Ser Trp Tyr Leu Asn His Ser Gly Asp Leu 425 430 420 Asn His Leu Pro Gly His Thr Phe Ala Ala Gln Gln Gln Thr Phe Pro 440 445 Asn Val Arg Glu Met Phe Asn Ser His Arg Leu Gly Ile Glu Asn Ser 455 460 Thr Leu Gly Glu Ser Gln Val Ser Gly Asn Ala Ser Cys Gln Leu Pro 470 475 Tyr Arg Ser Thr Pro Pro Leu Tyr Arg His Ala Ala Pro Tyr Ser Tyr 490 485 Asp Cys Thr Lys Tyr 500

<210> 3 <211> 4158 <212> DNA <213> Homo sapiens <220>

<220> <221> CDS <222> (187)..(1437)

<400> 3
cctttggctt tgaattgatc aggagacaaa gataatgcat ctacattttc gtcttctgtt 60
cttttattgg aaataagtgg cacgccccat tgccttctag tcgcctcccc gaagcgaaga 120
ggccgaagcg aagaggcctg gtgggttgtc tcaacatcct tttgctgaga atcgaatacg 180
cagccg atg aac agc cag gaa ggg tgc aag gaa acc ttg aac ggc atc 228
Met Asn Ser Gln Glu Gly Cys Lys Glu Thr Leu Asn Gly Ile

tac Tyr 15	cag Gln	ttc Phe	atc Ile	atg Met	gac Asp 20	cgc Arg	ttc Phe	ccc Pro	ttc Phe	tac Tyr 25	cgg Arg	gag Glu	aac Asn	aag Lys	cag Gln 30	276
				agc Ser 35												324
				cgc Arg												372
				gac Asp												420
				cgc Arg												468
		-		ctc Leu	_		_		_				-		_	516
				cac His 115												564
				agc Ser												612
				ctg Leu												660
				acg Thr												708
				gcg Ala												756
				cga Arg 195												804
				Gly ggc												852
				gcc Ala												900
gcc Ala	999 Gly 240	gcc Ala	gcc Ala	ggg Gly	ggc Gly	tac Tyr 245	cag Gln	tgc Cys	agc Ser	atg Met	cga Arg 250	gcg Ala	atg Met	agc Ser	ctg Leu	948

+1

tac acc ggg gcc gag cgg ccg gcg cac atg tgc gtc ccg ccc gcc ctg 996  Tyr Thr Gly Ala Glu Arg Pro Ala His Met Cys Val Pro Pro Ala Leu 255 260 265 270	
gac gag gcc ctc tcg gac cac ccg agc ggc ccc acg tcg ccc ctg agc 1044 Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Thr Ser Pro Leu Ser 275 280 285	
gct ctc aac ctc gcc gcc ggc cag gag ggc gcg ctc gcc acg ggc 1092 Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Ala Ala Thr Gly 290 295 300	
cac cac cac cag cac cac ggc cac cac cac	
ccg ccg gct ccc cag ccc cag ccg acg ccg cag ccc ggg gcc gcc	
gcg cag gcg gcc tcc tgg tat ctc aac cac agc ggg gac ctg aac cac 1236 Ala Gln Ala Ala Ser Trp Tyr Leu Asn His Ser Gly Asp Leu Asn His 335 340 345 350	
ctc ccc ggc cac acg ttc gcg gcc cag cag caa act ttc ccc aac gtg 1284 Leu Pro Gly His Thr Phe Ala Ala Gln Gln Gln Thr Phe Pro Asn Val 355 360 365	
cgg gag atg ttc aac tcc cac cgg ctg ggg att gag aac tcg acc ctc 1332 Arg Glu Met Phe Asn Ser His Arg Leu Gly Ile Glu Asn Ser Thr Leu 370 375 380	
ggg gag tcc cag gtg agt ggc aat gcc agc tgc cag ctg ccc tac aga 1380 Gly Glu Ser Gln Val Ser Gly Asn Ala Ser Cys Gln Leu Pro Tyr Arg 385 390 395	
tee acg ccg cct ctc tat cgc cac gca gcc ccc tac tee tac gac tgc 1428 Ser Thr Pro Pro Leu Tyr Arg His Ala Ala Pro Tyr Ser Tyr Asp Cys 400 405 410	
acg aaa tac tgacgtgtcc cgggacctcc cctccccggc ccgctccggc Thr Lys Tyr 415	
ttegettece ageceegace caaccagaca attaagggge tgcagagacg caaaaaagaa 1537	
acaaaacatg tccaccaacc ttttctcaga cccgggagca gagagcgggc acgctagccc 1597	
ccagccgtct gtgaagagcg caggtaactt taattcgccg ccccgtttct gggatcccag 1657	
gaaacccctc caaagggacg cagcccaaca aaatgagtat tggtcttaaa atcccctcc 1717	
cctaccagga cggctgtgct gtgctcgacc tgagctttca aaagttaagt tatggaccca 1777	
aatcccatag cgagccccta gtgactttct gtaggggtcc ccataggtgt atgggggtct 1837	
ctatagataa tatatgtgct gtgtgtaatt ttaaatttct ccaaccgtgc tgtacaaatg 1897	

tgtggatttg taatcaggct attttgttgt tgttgttgtt gttcagagcc attaatataa 1957 tatttaaagt tgagttcact ggataagttt ttcatcttgc ccaaccattt ctaactgcca 2017 aattgaattc aagaaaccga tgtgggtttt gtttcctgta caattatgag atataattct 2077 ttttcccatt gtaggtcttt tacaaaacaa gaaaataatt tatttttttg ttggtggata 2137 aagaagtcaa gtatctgata ctttttattt acaaagtgtg atggttttgt atagtaggtt 2197 ccaccctgag tattcctaaa agaaaaaaaa aaaaaaagct taaaaaactct aacttcatct 2257 gtgtttgtct tacgtggtct taatcgttgt acttacctta aaataaaccc atgttgtttt 2317 ttctgcccaa agtttggaca gtgtgtttgt gttgttgcat tttttacaaa cgaggtgtgt 2377 ttgcaaaccc acctgctttg attatttttg ttacacaggt gggtatatgt gtagacacat 2437 aaaaacgacc agagaatagg agcacaccc tgctgtcttg tttagtgaca gaaaaaggct 2497 tttgattaat tttaaaatcc cactctagga ttttttcttt tcgagaaacc gcccagttgg 2557 agggggetgc etgaaggace ggaceatgag tttgccgtga tgcattttet taaatgcaea 2617 aaaacatgct aattgtcaaa acaaacagtg ccactccatc tcagtgtcca gccgtcccca 2677 gtttaggagg tgaaggaagg gaagaataaa catttcccgt ttgctaactg caacccaqqq 2737 tgagtcctgc tttcccccga ttttataaaa tttgagcctc tttgcctgct ttaatagttt 2797 tccagagaat ttgaactggg ccaatgaagg tctgaagggg acggattttc tagcgtttga 2857 tatecatece cettagegge cagateagag gggaatttea gaetttatta etteteaatg 2917 tcatgtctaa atctacaccc tcatcgcagt gaaaaatttt aaaacctcat tacccttcaa 2977 aaataattta tgatatttt agagttctaa attcaagttt ttcaatatgt taaataatag 3037 agattatttt ttgttttcaa tgttaatatc tcgtctttta catttttaat agtaacatag 3097 tttttgtgaa atgtagetga egaaatgget ttattateta tttcaatgge tgaagteeae 3157 cacteceetg etggeeteta tgtgtgaatt tggggaccaa agetteatea atteceacce 3217 caqcaggtga gctgtacctt gctaatgctg aagttctttg tgagcttaac gtttcaagac 3277 cagatgattt tgctaaaggt gattttgctt gatgcagtgg cgctgaacgt aacccgggtg 3337 tttttgtcgt gttgttttca acatggcact ttatctccac gctatgttga aatagaatta 3397 qqqqaaqctt aaaqcataat aattqtcccc acatqtqcaa cacaqactct ttcaatctqt 3457 ggccccagag gtggcacaca gttaagactt ggcggctgtc tcattctttt tcataatgtg 3517 egggtteeeg ggtgteeggg tgetagaett teageaggee eeaggeeaga egggetttgg 3577 ttgagtgaac aggaggagga agttaaggag gtaggggtgg ggagagaccc tctccaagct 3637 gcagaagaag gtggcccaag ctccttgcct gcgtctgccg tgatggtttc attttacttc 3697

. 1

tgetegette atgetattig ecceaggaga agaggagagt attecagaeg gtaagegage 3757
tggetttte eetteectag aegittaaa gaaatettie tgaaagettig eeeteategt 3817
aagetitgaa aeegitggig teetgitagi ggegaggget gagagaaaeg eggagaaata 3877
aaggagageg aeggitgge tgagageece eaggietget gitgaaaeta agetgggett 3937
tigeaeetti aggaageett titaaagaag teetgetigi tgggggeegg aageecaagi 3997
gagtgggeet tgitgaggit alegggagg giettiaeea eteetiggig aaegitgggea 4057
aeggggggat tgitatetgaa geittattea ggietiege ggeageagag tggagaaeea 4117
ggeeettagt gigtagegge etggggatti tgggaeteat e 4158

<210> 4

<211> 417

<212> PRT

<213> Homo sapiens

<400> 4

Met Asn Ser Gln Glu Gly Cys Lys Glu Thr Leu Asn Gly Ile Tyr Gln
1 5 10 15

Phe Ile Met Asp Arg Phe Pro Phe Tyr Arg Glu Asn Lys Gln Gly Trp 20 25 30

Gln Asn Ser Ile Arg His Asn Leu Ser Leu Asn Glu Cys Phe Val Lys 35 40 45

Val Pro Arg Asp Asp Lys Lys Pro Gly Lys Gly Ser Tyr Trp Thr Leu 50 55 60

Asp Pro Asp Ser Tyr Asn Met Phe Glu Asn Gly Ser Phe Leu Arg Arg 65 70 75 80

Arg Arg Phe Lys Lys Asp Val Ser Lys Glu Lys Glu Glu Arg 85 90 95

Ala His Leu Lys Glu Pro Pro Pro Ala Ala Ser Lys Gly Ala Pro Ala 100 105 110

Thr Pro His Leu Ala Asp Ala Pro Lys Glu Ala Glu Lys Lys Val Val

Ile Lys Ser Glu Ala Ala Ser Pro Ala Leu Pro Val Ile Thr Lys Val 130 135 140

Glu Thr Leu Ser Pro Glu Ser Ala Leu Gln Gly Ser Pro Arg Ser Ala 145 150 155 160

Ala Ser Thr Pro Ala Gly Ser Pro Asp Gly Ser Leu Pro Glu His His 165 170 175

Ala Ala Pro Asn Gly Leu Pro Gly Phe Ser Val Glu Asn Ile Met 180 185 . 1

Thr Leu Arg Thr Ser Pro Pro Gly Gly Glu Leu Ser Pro Gly Ala Gly Arg Ala Gly Leu Val Val Pro Pro Leu Ala Leu Pro Tyr Ala Ala Ala Pro Pro Ala Ala Tyr Gly Gln Pro Cys Ala Gln Gly Leu Glu Ala Gly 225 230 235 Ala Ala Gly Gly Tyr Gln Cys Ser Met Arg Ala Met Ser Leu Tyr Thr Gly Ala Glu Arg Pro Ala His Met Cys Val Pro Pro Ala Leu Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Thr Ser Pro Leu Ser Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Ala Ala Thr Gly His His His Gln His His Gly His His Pro Gln Ala Pro Pro Pro Pro Pro 310 315 Ala Pro Gln Pro Gln Pro Thr Pro Gln Pro Gly Ala Ala Ala Gln 330 Ala Ala Ser Trp Tyr Leu Asn His Ser Gly Asp Leu Asn His Leu Pro Gly His Thr Phe Ala Ala Gln Gln Gln Thr Phe Pro Asn Val Arg Glu Met Phe Asn Ser His Arg Leu Gly Ile Glu Asn Ser Thr Leu Gly Glu Ser Gln Val Ser Gly Asn Ala Ser Cys Gln Leu Pro Tyr Arg Ser Thr Pro Pro Leu Tyr Arg His Ala Ala Pro Tyr Ser Tyr Asp Cys Thr Lys 410

Tyr

<210> 5

<211> 6021

<212> DNA

<213> Mus musculus

<220>

<221> exon

<222> (1649)..(4348)

<300>

<308> GenBank/Y08222

<309> 1997-05-14 <300> <301> Miura, N <303> Genomics <304> 41 <306> 489-492 <307> 1997

<400> 5

ctcgagtcaa aggtagcaca cataaaacct attttgctgc ttcggtacgt caagcaatgc 60 cactaaagtt teeteaceeg ceaaagetga aacagtgagt tetaatetet caaageettt 120 tgccgaaaat ctaaaggggg tggggggcta tggtggtggc gtggggggg ggtcggagaa 180 gaagaaagac tgagacaaat gttttatctg tcgccttctt ccctacccaa ccggaccaac 240 aacttccaga aggttctgcg aggcatagag ccattccgta gggacatctc ggtgcttctg 300 aggaagegga eegageaggg ateegatgae gaetggagat gttgaaggaa taaataceag 360 tccacaaata aacaaactgt ccccgggatt cctagaggga aggagcacgc ttgaaggtcg 420 gggaactccg agtcgctgtg cgtcaaggtt ggcataaaat taaaaaaaaa aaaagtcctt 480 cagttaccaq qecetetaaq gageeeetgg teeteagete acettateaa aactcagtaa 540 aacaaacago otgaaataca gtoaatttao aggatoocaa agatgotgao ogoggagtgg 600 gacccaegec gggccccggc aacagetagg gaagegggte egaggetaca cagtgeegeg 660 ctcctttgcg tttccagtga cgaagccggc gatggagtgc aggcttggag ctccccacgc 720 cgaacgggga caccagetee egggggetgg etgeettgte etaaceteca gacagegett 780 tcatagqtqq ggagaaggga gaggccggga tggatggcag ggaaagctag ccctcgtcta 840 tgcgggagag gagaccagga aagcaacagt tgggttcacg cgcttccctg aaccccacga 900 aattgtttgg aggacteaga tggateacet aagtageage gaagaegaag gaccaatggt 960 teettaggtg ttacetteec agtttggcat teecactaag cetteectee cageeegaee 1020 ccgtcgtgaa ggggagagga accgaattct ccaacccggc ctcctttgtg ggctcttcct 1080 caacctggaa gcgtcctgtg aattatccat cactgcattc aacaggccct acacgctcag 1140 tecgtttget etgaacecat tacaactagg eeeegataat taagaaatet aattattege 1200 ctcttcatcc attaataata ataaaaaaaa aatctccagg ctctttccta cttacaaggt 1260 cttgggggca aatctctgcc caacttcatc aattcgatgt tatatttcaa actaaacttc 1320 tttttatttt ccaaaggaac agggttttta atttttgete tggacacgtg gtetegttaa 1380 acaaaatgtg ataataaaat aaaattttat aagatgtaac tcatttttaa aagtcctcaa 1440 gttaacttga gctggggggg ggggagatct ggctaagagc atctgggtct tagagccgac 1500 ggattcaqqc gctcctcgtt ttgattggtg ccatccttct cgcagctgcc agatgattgg 1560 tgcaaacttc ctggaggggg cgcggcctga agaaagtaaa aactcgcttt gagccagaag 1620 acttttgaaa cttttcccaa tccctaaaag ggactttgct tctttttccg ggctcggccg 1680 cgcagcctct ccggacccta gctcgctgac gctgcgggct gcagttctcc tggcggggcc 1740 egagageege tgteteettt tetageacte ggaagggetg gtgtegetee aeggtegege 1800 gtggcgtctg tgccgccagc tcagggctgc cacccgccaa gccgagagtg cgcggccagc 1860 ggggccgcct gccgtgcacc cttcaggatg ccgatccgcc cggtcggctg aacccgagcg 1920 coggogtett cogogogtgg accordage tgccccqagt cggggctgcc tgcatcgctc 1980 egtecettee tgeteteetg eteegggeet egetegeege gggeegeagt eggtgegege 2040 aggeggegac egggegtetg ggaegeagea tgeaggegeg ttacteggta teggaeceea 2100 acgccctggg agtggtaccc tatttgagtg agcaaaacta ctaccgggcg gccggcagct 2160 acggcggcat ggccagcccc atgggcgtct actccggcca cccggagcag tacggcgccg 2220 gcatgggccg ctcctacgcg ccctaccacc accagcccgc ggcgcccaag gacctggtga 2280 agoogoocta cagotatata gogotoatoa coatggogat coagaacgog coagagaaga 2340 agateactet gaacggeate taccagttea teatggaceg ttteccette taccgegaga 2400 acaagcaggg ctggcagaac agcatccgcc acaacctgtc actcaatgag tgcttcgtga 2460 aagtqccqcq cqacqacaaq aagccgggca agggcagcta ctggacgctc gacccggact 2520 cctacaacat gttcgagaat ggcagettee tgcggcggcg gcggcgctte aagaagaagg 2580 atgtgcccaa ggacaaggag gagcgggccc acctcaagga gccgccctcg accacggcca 2640 agggegetee gacagggace eeggtagetg aegggeecaa ggaggeegag aagaaagteg 2700 tggttaagag cgaggcggcg tcccccgcgc tgccggtcat caccaaggtg gagacgctga 2760 gccccgaggg agcgctgcag gccagtccgc gcagcgcatc ctccacgccc gcaggttccc 2820 cagacggete getgeeggag caecaegeeg eggegeetaa egggetgeee ggetteageg 2880 tggagaccat catgacgctg cgcacgtcgc ctccgggcgg cgatctgagc ccagcggccg 2940 : 1

egegegeegg cetggtggtg ceaeegetgg caetgeeata egeegeageg ceaeeegeeg 3000 cttacacgca gccgtgcgcg cagggcctgg aggctgcggg ctccgcgggc taccagtgca 3060 gtatgcgggc tatgagtctg tacaccgggg ccgagcggcc cgcgcacgtg tgcgttccgc 3120 cegegetgga egaggetetg teggaceaec egageggeec eggeteeeeg eteggegeee 3180 teaacetege agegggteag gagggeget tgggggeete gggteaeeae caccageate 3240 acqqccacct ccaccqcaq qcqccaccqc ccqccccqca qccccctccc qcqccqcaqc 3300 cogcoaccca agecacctco tagtatota accaegaga agacetaase caceteecca 3360 gccacacgtt tgcaacccaa cagcaaactt tccccaacgt ccgggagatg ttcaactcgc 3420 accggctagg actggacaac tcgtccctcg gggagtccca ggtgagcaat gcgagctgtc 3480 agetgeecta tegagetaeg cegteectet acegecaege agececetae tettaegaet 3540 gcaccaaata ctgaggctgt ccagtccgct ccagccccag gaccgcaccg gcttcgcctc 3600 ctccatggga accttetteg acggageege agaaagegae ggaaagegee ceteteteag 3660 aaccaggage agagagetee gtgcaacteg caggtaactt atecgcaget cagtttgaga 3720 tctcagcgag tccctctaag ggggatgcag cccagcaaaa cgaaatacag atttttttt 3780 taatteette eectaceeag atgetgegee tgeteeettg gggetteata gattagetta 3840 tggaccaaac ccatagggac ccctaatgac ttctgtggag attctccacg ggcgcaagag 3900 gtctctccgg ataaggtgcc ttctgtaaac gagtgcggat ttgtaaccag gctattttgt 3960 tettgeccag ageetttaat ataatattta aagttgtgte caetggataa ggtttegtet 4020 tgcccaactg ttactgccaa attgaattca agaaacgtgt gtgggtcttt tctccccacg 4080 tcaccatgat aaaataggtc cctccccaaa ctgtaggtct tttacaaaaac aagaaaataa 4140 tttatttttt tgttgttgtt ggataacgaa attaagtatc ggatactttt aatttaggaa 4200 gtgcatggct tigtacagta gatgccatct ggggtattcc aaaaacacac caaaagactt 4260 taaaatttca atctcacctg tgtttgtctt atgtgatctc agtgttgtat ttaccttaaa 4320 ataaacccqt qttqtttttc tqcccaaaqt tcqqacaqaq tctttqtqtt cttqaatttt 4380 aaaagggaaa ttgtagtaag ccagttgtga ttgatttttg tgatgcaggt tggcctggta 4440 acgtggatgc atatacaggt tacaggacga tggagctetc gattagtaat agaagggget 4500 cttgatttgt tgaactatcc cgtcctgaga tatttttgtt ttctgctcga ggtaatctga 4560 gaaactgttc tccatccaca cacggacagg gctgcctgag ggcaacgtcc tgctggcctg 4620 ttaacgaaat gctttgcggg atgcagaaaa ctgttgccaa ttgtcaaaac aaaatggtgt 4680 caccetytet eggtytecay etyteetety ttagagggga gaaacegaga aaggacaaac 4740 ggcctgcagc ttgctaacct cagcgtagca ggagcctggg tgagtgctcg gctccctcca 4800 tttccttaga tgcggacttg ttgcccctgt tggcgtttta agagtgccag caagaagcaa 4860 agagggttgg taggtctctg gtatttaact gccggctttg ggatcagatt agaagtgaat 4920 ttcagtctga tttatttctt aatttgggct ttaaatattt tactccggcg tggtggaaaa 4980 agaagccact gtgcgcctcc agcatgatat tttagcgctg aaatggctct ggttttcagc 5040 aaaaaggaaa tagatcggga caaactctct aaaatgtacc tggctggctg gggtggggtc 5160 cttaccaatc tgctgcctga aagatacagc ttcagcacag gcctgcgtgt tggactttag 5220 gcatatcatg gattcccacg ccagttggta acctggactg tgctaatgga agttctttct 5280 gcacagaaca tgtaggccag gaggaggcag ggacccggga ggggggtgga ctttgcaggt 5340 catctgctta gcttagtggt ggccacgggt taacacgtat atagtgttac tgtttgaaac 5400 tecaagtttt atatetgtge tgttttgatg tagaatttgg ggaggtteet gatgataeta 5460 ccctacccgt gtatgtaaga cagtctttca acctgcagtg ccagaatgtg acccacactt 5520 cagtatette cataaagtgg ggggactaag aactggacag gggtgetgtg gaggggggca 5590 ggccaggtgt atcttggttc ctgagcagag cagagagett aggaaggggt cgggagatet 5640 etggtteete eeaacaetgg ttteattttg catggetete tteaaacete ttgeeceagg 5700 agaagcgagc tttgtccaag ccagctggct cgctcctttc ccagatgttt taggggcctc 5760 cctgaaagct tgccctcctc ttaagattca gaactcctga cccagggaaa gataggaggc 5820 tttgtggatg ggagcttttt tttaaagagg accgttctcg ttctcaagta ggtagctaga 5880 gagaagcccc ctggagcagg ccctacttgt gactgtcagg gaacccaggt tgtgttgtag 5940 gettttecca ggeeteecag ageageggtg tgaaaaaatg eggteetggg aaaagttggt 6000 ctggggtgtt gcttcctcga g 6021

<210> 6 <211> 2712 <212> DNA <213> Mus musculus

<220> <221> CDS <222> (422)..(1906) <300> <308> GenBank/Y08222 <309> 1997-05-14 <300> <301> Miura, N <303> Genomics <304> 41 <306> 489-492 <307> 1997 <400> 6 agggaetttg ettetttte egggetegge egegeageet eteeggaece tagetegetg 60 acqctqcqqq ctqcaqttct cctqqcqqqq cccqaqaqcc qctqtctcct tttctaqcac 120 teggaaggge tggtgteget eeacggtege gegtggegte tgtgeegeea geteaggget 180 gecaceegee aageegagag tgegeggeea geggggeege etgeegtgea eeetteagga 240 tgccgatccg cccggtcggc tgaacccgag cgccggcgtc ttccgcgcgt ggaccgcgag 300 getgeecega gteggggetg cetgeatege teegteectt eetgetetee tgeteeggge 360 ctegetegee gegggeegea gteggtgege geaggeggeg acegggegte tgggaegeag 420 c atg cag gcg cgt tac tcg gta tcg gac ccc aac gcc ctg gga gtg gta 469 Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala Leu Gly Val Val ccc tat ttg agt gag caa aac tac tac egg geg gee gge age tac gge Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala Gly Ser Tyr Gly 20 qqc atq qcc aqc ccc atq qqc qtc tac tcc qqc cac ccg qaq caq tac 565 Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His Pro Glu Gln Tyr gge gee gge atg gge ege tee tae geg eee tae eac eac eag eee geg 613 Gly Ala Gly Met Gly Arg Ser Tyr Ala Pro Tyr His His Gln Pro Ala gcg ccc aag gac ctg gtg aag ccg ccc tac agc tat ata gcg ctc atc 661 Ala Pro Lys Asp Leu Val Lys Pro Pro Tyr Ser Tyr Ile Ala Leu Ile 70 709 acc atg gcg atc cag aac gcg cca gag aag aag atc act ctg aac ggc Thr Met Ala Ile Gin Asn Ala Pro Glu Lys Lys Ile Thr Leu Asn Gly atc tac cag ttc atc atg gac cgt ttc ccc ttc tac cgc gag aac aag 757 Ile Tyr Gln Phe Ile Met Asp Arg Phe Pro Phe Tyr Arg Glu Asn Lys

105

.1

	g ggc n Gly															805
	gtg Val 130															853
	acg Thr															901
	g cgg n Arg															949
	gag Glu		_			_		_		_		_	-	_		997
-	ccg Pro				_	_	_	-			_		_		_	1045
aaa Lys	gtc Val 210	gtg Val	gtt Val	aag Lys	agc Ser	gag Glu 215	gcg Ala	gcg Ala	tcc Ser	ccc Pro	gcg Ala 220	ctg Leu	ccg Pro	gtc Val	atc Ile	1093
	aag Lys															1141
	agc Ser															1189
	cac His															1237
	atc Ile															1285
	gcc Ala 290															1333
	gca Ala															1381
	gct Ala															1429
	tac Tyr															1477

ctg gac gag gct ctg tcg gac cac ccg agc ggc ccc ggc tcc ccg ctc 152 Leu Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Gly Ser Pro Leu 355 360 365	<u>?</u> 5
ggc gcc ctc aac ctc gca gcg ggt cag gag ggc gcg ttg ggg gcc tcg 157 Gly Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Gly Ala Ser 370 375 380	3
ggt cac cac cac cag cat cac ggc cac ctc cac ccg cag gcg cca ccg 162 Gly His His Gln His His Gly His Leu His Pro Gln Ala Pro Pro 385 390 395 400	1
ccc gcc ccg cag ccc cct ccc gcg ccg cag ccc gcc acc cag gcc acc 166 Pro Ala Pro Gln Pro Pro Pro Ala Pro Gln Pro Ala Thr Gln Ala Thr 405 410 415	;9
tcc tgg tat ctg aac cac ggc ggg gac ctg agc cac ctc ccc ggc cac 171 Ser Trp Tyr Leu Asn His Gly Gly Asp Leu Ser His Leu Pro Gly His 420 425 430	.7
acg ttt gca acc caa cag caa act ttc ccc aac gtc cgg gag atg ttc 176 Thr Phe Ala Thr Gln Gln Gln Thr Phe Pro Asn Val Arg Glu Met Phe 435 440 445	i5
aac tcg cac cgg cta gga ctg gac aac tcg tcc ctc ggg gag tcc cag 181 Asn Ser His Arg Leu Gly Leu Asp Asn Ser Ser Leu Gly Glu Ser Gln 450 455 460	.3
gtg agc aat gcg agc tgt cag ctg ccc tat cga gct acg ccg tcc ctc 186 Val Ser Asn Ala Ser Cys Gln Leu Pro Tyr Arg Ala Thr Pro Ser Leu 465 470 475 480	1
tac ege cac gea gee eee tac tet tac gac tge ace aaa tac tga 190 Tyr Arg His Ala Ala Pro Tyr Ser Tyr Asp Cys Thr Lys Tyr 485 490 495	6
ggctgtccag teegetecag eeceaggace geaceggett egeeteetee atgggaacet 196	6
tettegaegg ageegeagaa agegegegeeete teteagaace aggageagag 202	6
ageteegtge aactegeagg taacttatee geageteagt ttgagatete agegagteee 208	6
tctaaggggg atgcagccca gcaaaacgaa atacagattt tttttttaat tccttcccct 214	6
acccagatge tgegeetget eeettgggge tteatagatt agettatgga eeaaacceat 220	6
agggacccct aatgacttct gtggagattc tccacgggcg caagaggtct ctccggataa 226	
ggtgccttct gtaaacgagt gcggatttgt aaccaggcta ttttgttctt gcccagagcc 232	
tttaatataa tatttaaagt tgtgtccact ggataaggtt tcgtcttgcc caactgttac 238	
tgccaaattg aattcaagaa acgtgtgtgg gtcttttctc cccacgtcac catgataaaa 244	
taggtccctc cccaaactgt aggtctttta caaaacaaga aaataattta tttttttgtt 250	
gttgttggat aacgaaatta agtatcggat acttttaatt taggaagtgc atggctttgt 256	6

acagtagatg ccatctgggg tattccaaaa acacaccaaa agactttaaa atttcaatct 2626 cacctgtgtt tgtcttatgt gatctcagtg ttgtatttac cttaaaataa acccgtgttg 2686 tttttctgcc caaaaaaaaa aaaaaa 2712

<210> 7 <211> 494 <212> PRT <213> Mus musculus

<400> 7 Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala Leu Gly Val Val 10 Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala Gly Ser Tyr Gly 20 25 Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His Pro Glu Gln Tyr 40 Gly Ala Gly Met Gly Arg Ser Tyr Ala Pro Tyr His His Gln Pro Ala 55 60 Ala Pro Lys Asp Leu Val Lys Pro Pro Tyr Ser Tyr Ile Ala Leu Ile 70 75 Thr Met Ala Ile Gln Asn Ala Pro Glu Lys Lys Ile Thr Leu Asn Gly 90 85 Ile Tyr Gln Phe Ile Met Asp Arg Phe Pro Phe Tyr Arg Glu Asn Lys 100 105 110 Gln Gly Trp Gln Asn Ser Ile Arg His Asn Leu Ser Leu Asn Glu Cys 115 120 Phe Val Lys Val Pro Arg Asp Asp Lys Lys Pro Gly Lys Gly Ser Tyr 135 140 Trp Thr Leu Asp Pro Asp Ser Tyr Asn Met Phe Glu Asn Gly Ser Phe 150 155 Leu Arg Arg Arg Arg Phe Lys Lys Lys Asp Val Pro Lys Asp Lys 175 170 165 Glu Glu Arg Ala His Leu Lys Glu Pro Pro Ser Thr Thr Ala Lys Gly 185 Ala Pro Thr Gly Thr Pro Val Ala Asp Gly Pro Lys Glu Ala Glu Lys 200 205 Lys Val Val Val Lys Ser Glu Ala Ala Ser Pro Ala Leu Pro Val Ile 215 220 Thr Lys Val Glu Thr Leu Ser Pro Glu Gly Ala Leu Gln Ala Ser Pro 230 235 Arg Ser Ala Ser Ser Thr Pro Ala Gly Ser Pro Asp Gly Ser Leu Pro 245 250 255 Glu His His Ala Ala Ala Pro Asn Gly Leu Pro Gly Phe Ser Val Glu 265 270 260 Thr Ile Met Thr Leu Arg Thr Ser Pro Pro Gly Gly Asp Leu Ser Pro 280 285 Ala Ala Arg Ala Gly Leu Val Val Pro Pro Leu Ala Leu Pro Tyr 295 300 Ala Ala Ala Pro Pro Ala Ala Tyr Thr Gln Pro Cys Ala Gln Gly Leu 305 310 315 Glu Ala Ala Gly Ser Ala Gly Tyr Gln Cys Ser Met Arg Ala Met Ser 330 Leu Tyr Thr Gly Ala Glu Arg Pro Ala His Val Cys Val Pro Pro Ala 340 345 350 Leu Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Gly Ser Pro Leu

360

365

::::

·::::

<210> 8

```
Gly Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Gly Ala Ser
                        375
Gly His His His Gln His His Gly His Leu His Pro Gln Ala Pro Pro
                                     . 395
385
                    390
Pro Ala Pro Gln Pro Pro Pro Ala Pro Gln Pro Ala Thr Gln Ala Thr
                                    410
                405
Ser Trp Tyr Leu Asn His Gly Gly Asp Leu Ser His Leu Pro Gly His
                                425
                                                    430
Thr Phe Ala Thr Gln Gln Gln Thr Phe Pro Asn Val Arg Glu Met Phe
                            440
                                                445
Asn Ser His Arg Leu Gly Leu Asp Asn Ser Ser Leu Gly Glu Ser Gln
                        455
                                            460
Val Ser Asn Ala Ser Cys Gln Leu Pro Tyr Arg Ala Thr Pro Ser Leu
                                        475
                    470
Tyr Arg His Ala Ala Pro Tyr Ser Tyr Asp Cys Thr Lys Tyr
                                    490
```

П

```
<211> 3289
<212> DNA
<213> Homo sapiens
<300>
<301> Miura, N
<303> Genomics
<304> 41
<306> 489-492
<307> 1997
<308> GenBank/Y08223
<309> 1997-05-14
gaattoggag gattaagttg toagtoagca ogttgetace ttoccetota tgcactoogc 60
tgcctggctc ctcggcgggg agcgagggaa actcagtttg tagggtttac ctctaaaacc 120
tegataggtt atcettgaeg acceegagee tggaaactee etgttgatga ttaattattt 180
gattaaataa gtataacatc caggagaggc cctgccattc caatccagcg cgtttgcttt 240
tgaatccatt acacctgggc ccccataatt aggaaatcta attattcgct tcatcactca 300
ttaataagaa aaatgtccca ggatcattgc tacttacaag gtctttggga gagatatttt 360
actictattaa ticattictat tittatattic aaattgatti tittitaacag aggaaagtgg 420
ctatcttttt gttttgggca tgtgggccca ttcaccaaaa tgtgatcata aaataaattt 480
taataagata taacttttta aaaagttttc aagtgaagac ggagtcgccg cggaggccgg 540
ggcggcgggg tcttagagcc gacggattcc tgcgctcctc gccccgattg gcgccggact 600
cctctcagct gccgggtgat tggctcaaag ttccgggagg gggcgtggcc cgaggaaagt 660
aaaaactcgc tttcagcaag aagacttttg aaacttttcc caatccctaa aagggacttg 720
geetetttt etgggeteag eggggeagee geteggaeee eggegegetg accetegggg 780
ctgccgattc gctgggggct tggagagcct cctgcgcccc tcctcgcgcg ggccgagggt 840
ccaccttggt ccccaggccg cggcgtctcc gctgggtccg cggccgcccg cctgcccgcg 900
ctgccgccgc cgggtcctgg agccagcgag gagcggggcc ggcgctgcgc ttgcccgggg 960
cgcgccctcc aggatgccga tccgcccggt ccgctgaaag cgcgccccc tgctcggccc 1020
gagegaegae gaeegegeae eetegeeeeg gaggetgeea ggagaeeggg geegeeeete 1080
ccgctccct cctctcccc tctggctctc tcgcgctctc tcgctctcag ggcccccctc 1140
gctcccccgg ccgcagtccg tgcgcgaggg cgccggcgag ccgtctcgga agcagcatgc 1200
aggegegeta etecgtgtee gaccecaacg ceetgggagt ggtgeeetae etgagegage 1260
```

agaattacta cegggetgeg ggeagetacg geggeatgge cageeceatg ggegtetatt 1320 ceggecacee ggageagtac agegegggga tgggeegete etacgegece taccaccace 1380

```
accagecege ggegeetaag gacetggtga ageegeeeta cagetacate gegeteatea 1440
ccatggccat ccagaacgcg cccgagaaga agatcacctt gaacggcatc taccagttca 1500
tcatggaccg cttccccttc taccgggaga acaagcaggg ctggcagaac agcatccgcc 1560
acaacctete geteaacgag tgettegtea aggtgeeeeg egaegacaag aageceggea 1620
agggcagtta ctggaccctg gacccggact cctacaacat gttcgagaac ggcagcttcc 1680
tgcggcgccg gcggcgcttc aaaaagaagg acgtgtccaa ggagaaggag gagcgggccc 1740
acctcaagga geegeeeeg geggegteea agggegeeee ggeeaeeeee cacctagegg 1800
acgcccccaa ggaggccgag aagaaggtgg tgatcaagag cgaggcggcg tccccggcgc 1860
tgccggtcat caccaaggtg gagacgctga gccccgagag cgcgctgcag ggcagcccgc 1920
geagegege etecaegee geeggeteec cegaeggtte getgeeggag caecaegeeg 1980
cggcgcccaa cgggctgcct ggcttcagcg tggagaacat catgaccctg cgaacgtcgc 2040
egeegggegg agagetgage eegggggeeg gaegegeggg cetggtggtg eegeegetgg 2100
cgctgccata cgccgccgcg ccgcccgccg cctacggcca gccgtgcgct cagggcctgg 2160
aggccggggc cgccgggggc taccagtgca gcatgcgagc gatgagcctg tacaccgggg 2220
ccgageggee ggegeacatg tgegteeege cegeectgga egaggeeete teggaecace 2280
cgagcggccc cacgtcgccc ctgagcgctc tcaacctcgc cgccggccag gagggcgcgc 2340
tegeegeeae gggeeaeeae caceageaee aeggeeaeea eeaeeegeag gegeegeege 2400
coccegecage tecceages cagecages egeageseg georgesege gegeages 2460
cctcctggta tctcaaccac agoggggacc tgaaccacct ccccggccac acgttcgcgg 2520
cccagcagca aactttcccc aacgtgcggg agatgttcaa ctcccaccgg ctggggattg 2580
aqaactcgac cctcggggag tcccaggtga gtggcaatgc cagctgccag ctgccctaca 2640
gatecaegee geetetetat egecaegeag eccectaete etaegaetge aegaaataet 2700
gacgtgtccc gggacctccc ctccccggcc cgctccggct tegettccca gccccgaccc 2760
aaccagacaa ttaaggggct gcagagacgc aaaaaagaaa caaaacatgt ccaccaacct 2820
tttctcagac cogggagcag agagogggca cgctagcocc cagoogtotg tgaagagogo 2880
aggtaacttt aattegeege eeegtttetg ggateeeagg aaaceeetee aaagggaege 2940
agoccaacaa aatgagtatt ggtottaaaa toccootcoo ctaccaggac ggotgtgotg 3000
tgctcgacct gagctttcaa aagttaagtt atggacccaa atcccatagc gagcccctag 3060
tgactttetg taggggteec cataggtgta tgggggtete tatagataat atatgtgetg 3120
tgtgtaattt taaatttoto caacogtgot gtacaaatgt gtggatttgt aatcaggota 3180
ttttgttgtt gttgttgttg ttcagagcca ttaatataat atttaaagtt gagttcactg 3240
gataagtttt tcatcttgcc caaccatttc taactgccaa attgaattc
```

```
<210> 9
<211> 1506
<212> DNA
<213> Homo sapiens
<220>
<221> CDS
<222> (1)..(1506)
<300>
<308> GenBank/NM 005251
<309> 1999-12-23
<400> 9
atg cag gcg cgc tac tcc gtg tcc gac ccc aac gcc ctg gga gtg gtg
Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala Leu Gly Val Val
ccc tac ctg agc gag cag aat tac tac cgg gct gcg ggc agc tac ggc
                                                                   96
Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala Gly Ser Tyr Gly
ggc atg gcc agc ccc atg ggc gtc tat tcc ggc cac ccg gag cag tac
                                                                   144
Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His Pro Glu Gln Tyr
                             40
```

				tcc Ser 55								192
				gtg Val								240
				aac Asn								288
				atg Met								336
				agc Ser								384
-	_	_		cgc Arg 135	-	-	_	_		_	 _	432
				gac Asp								480
				cgc Arg								528
				ctc Leu								576
				cac His								624
				agc Ser 215								672
	_		 _	ctg Leu	-			-	 _	-	 -	720
				acg Thr								768
				gcg Ala								816

	aac Asn												864
	ggg Gly 290												912
	gcc Ala												960
	gag Glu												1008
	ctg Leu												1056
-	ctg Leu	_	 -		•	-		-	_		_	_	1104
	agc Ser 370												1152
	ggc Gly												1200
	ccc Pro												1248
	gcg Ala												1296
	cac His												1344
	gtg Val 450												1392
	ctc Leu												1440
	aga Arg						Arg						1488
	tgc Cys			tga									1506

<210> 10 <211> 501 <212> PRT <213> Homo sapiens

<400> 10 Met Gln Ala Arg Tyr Ser Val Ser Asp Pro Asn Ala Leu Gly Val Val 10 Pro Tyr Leu Ser Glu Gln Asn Tyr Tyr Arg Ala Ala Gly Ser Tyr Gly 25 Gly Met Ala Ser Pro Met Gly Val Tyr Ser Gly His Pro Glu Gln Tyr 35 40 Ser Ala Gly Met Gly Arg Ser Tyr Ala Pro Tyr His His His Gln Pro 55 Ala Ala Pro Lys Asp Leu Val Lys Pro Pro Tyr Ser Tyr Ile Ala Leu 70 Ile Thr Met Ala Ile Gln Asn Ala Pro Glu Lys Lys Ile Thr Leu Asn 85 90 Gly Ile Tyr Gln Phe Ile Met Asp Arg Phe Pro Phe Tyr Arg Glu Asn 105 Lys Gln Gly Trp Gln Asn Ser Ile Arg His Asn Leu Ser Leu Asn Glu 115 120 125 Cys Phe Val Lys Val Pro Arg Asp Asp Lys Lys Pro Gly Lys Gly Ser 135 Tyr Trp Thr Leu Asp Pro Asp Ser Tyr Asn Met Phe Glu Asn Gly Ser 150 155 Phe Leu Arg Arg Arg Arg Phe Lys Lys Asp Val Ser Lys Glu 170 Lys Glu Glu Arg Ala His Leu Lys Glu Pro Pro Pro Ala Ala Ser Lys 185 190 180 Gly Ala Pro Ala Thr Pro His Leu Ala Asp Ala Pro Lys Glu Ala Glu 200 205 Lys Lys Val Val Ile Lys Ser Glu Ala Ala Ser Pro Ala Leu Pro Val 215 220 Ile Thr Lys Val Glu Thr Leu Ser Pro Glu Ser Ala Leu Gln Gly Ser 230 235 Pro Arg Ser Ala Ala Ser Thr Pro Ala Gly Ser Pro Asp Gly Ser Leu 245 250 Pro Glu His His Ala Ala Pro Asn Gly Leu Pro Gly Phe Ser Val 265 270 260 Glu Asn Ile Met Thr Leu Arg Thr Ser Pro Pro Gly Gly Glu Leu Ser 280 Pro Gly Ala Gly Arg Ala Gly Leu Val Val Pro Pro Leu Ala Leu Pro 290 295 300 Tyr Ala Ala Ala Pro Pro Ala Ala Tyr Gly Gln Pro Cys Ala Gln Gly 310 315 Leu Glu Ala Gly Ala Ala Gly Gly Tyr Gln Cys Ser Met Arg Ala Met 325 330 Ser Leu Tyr Thr Gly Ala Glu Arg Pro Ala His Met Cys Val Pro Pro 345 Ala Leu Asp Glu Ala Leu Ser Asp His Pro Ser Gly Pro Thr Ser Pro 355 360 Leu Ser Ala Leu Asn Leu Ala Ala Gly Gln Glu Gly Ala Leu Ala Ala 375 380 Thr Gly His His Gln His His Gly His His Pro Gln Ala Pro 390 395

```
Pro Pro Pro Pro Ala Pro Gln Pro Gln Pro Thr Pro Gln Pro Gly Ala
                                     410
Ala Ala Ala Gln Ala Ala Ser Trp Tyr Leu Asn His Ser Gly Asp Leu
                                                     430
            420
                                425
Asn His Leu Pro Gly His Thr Phe Ala Ala Gln Gln Gln Thr Phe Pro
                            440
                                                445
Asn Val Arg Glu Met Phe Asn Ser His Arg Leu Gly Ile Glu Asn Ser
                        455
                                            460
Thr Leu Gly Glu Ser Gln Val Ser Gly Asn Ala Ser Cys Gln Leu Pro
                    470
                                        475
Tyr Arg Ser Thr Pro Pro Leu Tyr Arg His Ala Ala Pro Tyr Ser Tyr
                485
                                    490
Asp Cys Thr Lys Tyr
            500
```

```
<210> 11
<211> 327
<212> DNA
<213> Homo sapiens
<300>
<308> GenBank/AW271272
<309> 2000-01-03
<300>
<301> Strausberg, Robert
<303> Trends Genet.
<304> 16
<305> 3
<306> 103-106
<307> 2000
<400> 11
ttttttttac attttcgtct tctgttcttg tgattggaaa taagtggcac gccccattgc 60
acateetttt getgagaate gaataegeag eegatgaaca geeaggaagg gtgeaaggaa 180
accttgaacg gcatctacca gttcatcatg gaccgcttcc ccttctaccg ggagaacaag 240
cagggctggc agaacagcat ccgccacaac ctctcgctca acgagtgctt cgtcaaggtg 300
ccccgcgacg acaagaagcc cggcaag
<210> 12
```

```
<212> DNA
<213> Homo sapiens

<300>
<308> GenBank/AW793237
<309> 2000-05-16


<300>
<301> Dias Neto, E
<303> Proc. Natl. Acad. Sci. U.S.A.
<304> 97
<306> 3491-3496
<307> 2000
```


<211> 147


<400> 12
ccgtctgaga atcgaatacg cagccgatga acagccagga agggtgcaag gaaaccttga 60
acggcatcta ccagttcatc atggaccgct tececttcta ccgggagaac aagcagggct 120
ggcagaacag catccgccac aacctct 147

Transcription Start Site

Fig. 1







## Fig. 3

E-box-like	HUMAN AAGGGTGCAAGGAAACCTGAAATACAAATGTTCTCCCTGAAGCCCTTCCCTGCCCAAC	CGGAGAAGAAGACTGAGACAAATGTTTTATCTGTCGCCTTCTTCCTACCCAAC
200 Splice E-	AAGGGTGCAAGGAAACCTGAA	CGGAGAAGAAGAAGACTGAG
• •	HUMAN	MOUSE

HUMAN CAGACCAGCAACTICCAAAATICIGCCGIGTITAGCCTIGTIAAAGGGGGIGTCICA	CGGACCAACAACTTCCAGAAGGTTCTGCGAGGCATAGAGCCATTCCGTAGGGACATCTCG
CAGACCAGCA	CGGACCAACA
HUMAN	MOUSE

	FOT VIIGAL TING
HUMAN	CTCCTTCAGGGAAAGTGGGAAAAGGGGATCTGATTATTGAGGTGTGGAAGGAAT
MOUSE	GIGCTICTGAGGAAGCGGACCGAGCAGGGATCCGATGACGACTGGAGATGTTGAAGGAAT

Ets-like	GGGATTCCTAGAGGGAAGGAGAATC	CCGGGATTCCTAGAGGGAAGGAGCACGC
Forkhead-like	AAATAATCAGTCCACAAATAAACAAACTGTCCGGGATTCCTAGAGGGAAGGAAATC	AAATA-CCAGTCCACAAATAAACAAACTGTCCCCGGGATTCCTAGAGGAAGGA
	HUMAN	MOUSE

HUMAN CTTGAAGGAGATCCAAGTCGCTCCAGGTCTGCCTGCCGAATAATATC

(1250)	N TGCCATTCCAATC-CAGCGCGTTTGCTTTTGAATCCATTACACCTGGGCCCCCATAATTA SE GCCCTACACGCTCAGTCCGTTTGCTC-TGAACCCATTACAACTAGGCCCCGATAATTA	NN GGAAATCTAATTATTCGCTTCATCACTCATTAATAAGAAAAATGTCCCAGGAT SE AGAAATCTAATTATTCGCCTCTTCATCCATTAATAATAAAAAAAA	NN CATTGCTACTTACAAGGTCTTTGGGAGAGATATTTTACTCTATTAATCCATTCTATTTA SE CTTTCCTACTTACAAGGTCTTGGGGGCAAATCTCTGCCCAACTTCATCAATTCGATGTTA	NN TATTTCAAATTGATTTTTTTTAACAGGAAAGTGGCTATCTTTTTGTTTTG	NN GGCATGTGGGCCCATTCACCAAATGTGATCATAAAATAAAATTTTAATAAGATATAACT~ SE GACACGTGGTCTCGTTAAACAAAATGTGATAATAAAATAAAATTTTATAAGATGTAACTC	NN -TTTTAAAAAGTTTTCAAGTGAAGACGGAGTCGCCGCGGGAGGCCGGGGGGGGGG	CGGGGTCTTAGAGCCGACGGATTCCTGCGCTCCTCGCCCCGATTGGC	NA CAGCTGCCGGGTGATTGGCTCAAAGTTCCGGGAGGGGGGGCGTGGCCCGAGGAAAGTAAAAA SE CAGCTGCCAGATGATTGGTGCAAACTTCCTGGAGGGGGGGG	
	HUMAN	HUMAN MOUSE	HUMAN	HUMAN MOUSE	HUMAN MOUSE	HUMAN	HUMAN	HUMAN	

CTCGCTTTCAGCAAGAAGACTTTTGAAACTTTTCCCAATCCCTAAAAGGGACTTGGCCTC- (1763) HUMAN MOUSE